

RSL24D1 shRNA (m) Lentiviral Particles: sc-141489-V

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. RSL24D1, also known as L30, RPL24L or HRP-L30-iso, is a 163 amino acid nuclear protein that shares a low level of similarity with Ribosomal Protein L24 (MRP-L24). Like other ribosomal proteins, RSL24D1 is involved in the biogenesis of the large 60S subunit and, during biogenesis, it is associated with nuclear and cytoplasmic pre-60S particles where it mediates proper protein docking. Once biogenesis is complete, RSL24D1 dissociates from the particles and is thought to be exchanged for Ribosomal Protein L24.

REFERENCES

1. Johnson, K.R. 1993. Characterization of cDNA clones encoding the human homologue of *Saccharomyces cerevisiae* Ribosomal Protein L30. *Gene* 123: 283-285.
2. Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T.J., Tanaka, T. and Page, D.C. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
3. Uechi, T., Tanaka, T. and Kenmochi, N. 2001. A complete map of the human ribosomal protein genes: assignment of 80 genes to the cytogenetic map and implications for human disorders. *Genomics* 72: 223-230.
4. Yoshihama, M., Uechi, T., Asakawa, S., Kawasaki, K., Kato, S., Higa, S., Maeda, N., Minoshima, S., Tanaka, T., Shimizu, N. and Kenmochi, N. 2002. The human ribosomal protein genes: sequencing and comparative analysis of 73 genes. *Genome Res.* 12: 379-390.

CHROMOSOMAL LOCATION

Genetic locus: Rsl24d1 (mouse) mapping to 9 D.

PRODUCT

RSL24D1 shRNA (m) Lentiviral Particles are concentrated, transduction-ready viral particles containing a target-specific construct that encodes a 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see RSL24D1 siRNA (m): sc-141489 and RSL24D1 shRNA Plasmid (m): sc-141489-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

RSL24D1 shRNA (m) Lentiviral Particles is recommended for the inhibition of RSL24D1 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

RSL24D1 (3H2): sc-100840 is recommended as a control antibody for monitoring of RSL24D1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RSL24D1 gene expression knockdown using RT-PCR Primer: RSL24D1 (m)-PR: sc-141489-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.