

BRD9 shRNA Plasmid (m): sc-141743-SH

BACKGROUND

The bromodomain is an approximately 70 amino acid region that is found in a variety of proteins and is thought to be involved in protein-protein interactions and transcriptional activation. BRD9 (bromodomain-containing protein 9), also known as PRO9856 or LAVS3040, is a 501 amino acid protein that contains one bromodomain and may be involved in various protein interactions throughout the cell. The gene encoding BRD9 maps to chromosome 5p15.33 and is expressed as four isoforms due to alternative splicing events. With 181 million base pairs encoding around 1,000 genes, chromosome 5 comprises about 6% of human genomic DNA. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm on chromosome 5 is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

1. Saltman, D.L., et al. 1993. A physical map of 15 loci on human chromosome 5q23-q33 by two-color fluorescence *in situ* hybridization. *Genomics* 16: 726-732.
2. Kadmon, M., et al. 2001. Duodenal adenomatosis in familial adenomatous polyposis coli. A review of the literature and results from the Heidelberg Polyposis Register. *Int. J. Colorectal Dis.* 16: 63-75.
3. South, S.T., et al. 2006. A new genomic mechanism leading to Cri-du-chat syndrome. *Am. J. Med. Genet. A* 140: 2714-2720.
4. Aretz, S., et al. 2007. Somatic APC mosaicism: a frequent cause of familial adenomatous polyposis (FAP). *Hum. Mutat.* 28: 985-992.
5. Cleaver, J.E., et al. 2007. Cockayne syndrome exhibits dysregulation of p21 and other gene products that may be independent of transcription-coupled repair. *Neuroscience* 145: 1300-1308.
6. Du, H.Y., et al. 2007. Telomerase reverse transcriptase haploinsufficiency and telomere length in individuals with 5p-syndrome. *Aging Cell* 6: 689-697.
7. Herry, A., et al. 2007. Redefining monosomy 5 by molecular cytogenetics in 23 patients with MDS/AML. *Eur. J. Haematol.* 78: 457-467.
8. Kang, J.U., et al. 2008. Gain at chromosomal region 5p15.33, containing TERT, is the most frequent genetic event in early stages of non-small cell lung cancer. *Cancer Genet. Cytogenet.* 182: 1-11.

CHROMOSOMAL LOCATION

Genetic locus: Brd9 (mouse) mapping to 13 C1.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PRODUCT

BRD9 shRNA Plasmid (m) is a pool of 3 target-specific lentiviral vector plasmids each encoding 19-25 nt (plus hairpin) shRNAs designed to knock down gene expression. Each plasmid contains a puromycin resistance gene for the selection of cells stably expressing shRNA. Each vial contains 20 µg of lyophilized shRNA plasmid DNA. Suitable for up to 20 transfections. Also see BRD9 siRNA (m): sc-141743 and BRD9 shRNA (m) Lentiviral Particles: sc-141743-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized shRNA plasmid DNA at 4° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at 4° C for short term storage or -80° C for long term storage. Avoid repeated freeze thaw cycles.

Resuspend lyophilized shRNA plasmid DNA in 200 µl of the deionized water provided. Resuspension of the shRNA plasmid DNA in 200 µl of deionized water makes a 0.1 µg/µl solution in a 10 mM Tris, 1 mM EDTA buffered solution.

APPLICATIONS

BRD9 shRNA Plasmid (m) is recommended for the inhibition of BRD9 expression in mouse cells.

SUPPORT REAGENTS

For optimal shRNA Plasmid transfection efficiency, Santa Cruz Biotechnology's shRNA Plasmid Transfection Reagent: sc-108061 (0.2 ml) and shRNA Plasmid Transfection Medium: sc-108062 (20 ml) are recommended. Control shRNAs are available as 20 µg lyophilized plasmid DNA. Each encodes a scrambled shRNA sequence that will not lead to the specific degradation of any known cellular mRNA. Control shRNA Plasmids include: sc-108060, sc-108065 and sc-108066.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor BRD9 gene expression knockdown using RT-PCR Primer: BRD9 (m)-PR: sc-141743-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.