

## Btnl6 siRNA (m): sc-141787

### BACKGROUND

Milk lipid globules of various species are surrounded by a membrane structure that is separated from the triglyceride core of the globule by a densely staining fuzzy coat layer of 10- to 50-nm thickness. This internal coat structure remains attached to the membrane during isolation and extraction with low- and high-salt buffers, is insoluble in nondenaturing detergents and is enriched by an acidic glycoprotein known as Butyrophilin. A major protein associated with fat droplets in the milk of many species, Butyrophilin is a member of the immunoglobulin superfamily, a large family of proteins comprised of components of the immune system, plasma membrane receptors and adhesive molecules. Butyrophilin is a glycoprotein that is specifically expressed on the apical surface of mammary epithelial cells during lactation and becomes incorporated as an integral protein into the membrane of the milk fat globule during the budding and secretion of fat droplets into milk. Btnl6 (butyrophilin-like 6), also known as EG624681, is a 539 amino acid protein containing one B30.2/SPRY domain. Btnl6 is encoded by a gene that maps to mouse chromosome 17 B1.

### REFERENCES

1. Franke, W.W., et al. 1981. Antibodies to the major insoluble milk fat globule membrane-associated protein: specific location in apical regions of lactating epithelial cells. *J. Cell Biol.* 89: 485-494.
2. Mather, I.H., et al. 1993. A review of the molecular and cellular biology of butyrophilin, the major protein of bovine milk fat globule membrane. *J. Dairy Sci.* 76: 3832-3850.
3. Vernet, C., et al. 1993. Evolutionary study of multigenic families mapping close to the human MHC class I region. *J. Mol. Evol.* 37: 600-612.
4. Ogg, S.L., et al. 1996. Structural organization and mammary-specific expression of the butyrophilin gene. *Mamm. Genome* 7: 900-905.
5. Tazi-Ahnini, R., et al. 1997. Cloning, localization, and structure of new members of the butyrophilin gene family in the juxta-telomeric region of the major histocompatibility complex. *Immunogenetics* 47: 55-63.
6. Rhodes, D.A., et al. 2001. The cluster of BTN genes in the extended major histocompatibility complex. *Genomics* 71: 351-362.
7. Robenek, H., et al. 2006. Butyrophilin controls milk fat globule secretion. *Proc. Natl. Acad. Sci. USA* 103: 10385-10390.

### CHROMOSOMAL LOCATION

Genetic locus: Btnl6 (mouse) mapping to 17 B1.

### PRODUCT

Btnl6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Btnl6 shRNA Plasmid (m): sc-141787-SH and Btnl6 shRNA (m) Lentiviral Particles: sc-141787-V as alternate gene silencing products.

For independent verification of Btnl6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141787A, sc-141787B and sc-141787C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

Btnl6 siRNA (m) is recommended for the inhibition of Btnl6 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Btnl6 gene expression knockdown using RT-PCR Primer: Btnl6 (m)-PR: sc-141787-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.