

LAMP-5 siRNA (m): sc-141851

BACKGROUND

Lysosome-associated membrane proteins (LAMP) are glycosylated type I membrane proteins that play a role in the biogenesis of the pigment melanin. LAMP-5, also known as C20orf103, is a 280 amino acid protein that belongs to the LAMP family and is mainly found in cytoplasmic vesicles of the neuronal cell body. LAMP-5 is up-regulated upon CpG activation and down-regulated upon Toll-like receptor (TLR) ligands.

REFERENCES

1. Febbraio, M., et al. 1990. Identification and characterization of LAMP-1 as an activation-dependent platelet surface glycoprotein. *J. Biol. Chem.* 265: 18531-18537.
2. Salopek, T.G., et al. 1996. Induction of melanogenesis during the various melanoma growth phases and the role of tyrosinase, lysosome-associated membrane proteins, and p90 calnexin in the melanogenesis cascade. *J. Invest. Dermatol. Symp. Proc.* 1: 195-202.
3. Wilke, S., et al. 2012. Crystal structure of the conserved domain of the DC lysosomal associated membrane protein: implications for the lysosomal glycolyx. *BMC Biol.* 10: 62.
4. Defays, A., et al. 2011. BAD-LAMP is a novel biomarker of nonactivated human plasmacytoid dendritic cells. *Blood* 118: 609-617.
5. Yang, K., et al. 2009. Immune responses to T-cell epitopes of SARS CoV-N protein are enhanced by N immunization with a chimera of lysosome-associated membrane protein. *Gene Ther.* 16: 1353-1362.

CHROMOSOMAL LOCATION

Genetic locus: Lamp5 (mouse) mapping to 2 F3.

PRODUCT

LAMP-5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LAMP-5 shRNA Plasmid (m): sc-141851-SH and LAMP-5 shRNA (m) Lentiviral Particles: sc-141851-V as alternate gene silencing products.

For independent verification of LAMP-5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141851A, sc-141851B and sc-141851C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LAMP-5 siRNA (m) is recommended for the inhibition of LAMP-5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

LAMP-5 (E-8): sc-398190 is recommended as a control antibody for monitoring of LAMP-5 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LAMP-5 gene expression knockdown using RT-PCR Primer: LAMP-5 (m)-PR: sc-141851-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.