

β3Gn-T9 siRNA (m): sc-141927

BACKGROUND

A family of human β 1,3-galactosyltransferases (β3Gn-Ts) consists of nine members (β3Gn-T1, -T2, -T3, -T4, -T5, -T6, -T7, -T8 and -T9). β3Gn-T1 catalyzes the formation of type 1 oligosaccharides. β3Gn-T2 converts lacto-N-triose II into lacto-N-tetraose and lacto-N-neotetraose and can form a heterodimer with β3Gn-T8, which, as a complex, exhibits higher enzymatic activity. Unlike the ubiquitously expressed β3Gn-T2, β3Gn-T3 is specifically expressed in colon, jejunum, stomach, esophagus, placenta and trachea, and β3Gn-T4 is mainly expressed in brain. β3Gn-T5 is essential for the biosynthesis of Lewis antigens and may play a role in gastric cancer as a result of its participation in chronic *H. pylori* infection. β3Gn-T6 may be a useful marker for distinguishing between benign adenomas and premalignant lesions. β3Gn-T7 acts as an anti-migration factor for a lung cancer cell line. β3Gn-T9 is a single-pass type II membrane protein localized to the Golgi apparatus membrane.

REFERENCES

1. Shiraishi, N., Natsume, A., Togayachi, A., Endo, T., Akashima, T., Yamada, Y., Imai, N., Nakagawa, S., Koizumi, S., Sekine, S., Narimatsu, H. and Sasaki, K. 2001. Identification and characterization of three novel β 1,3-N-acetylglucosaminyltransferases structurally related to the β 1,3-galactosyltransferase family. *J. Biol. Chem.* 276: 3498-3507.
2. Seko, A. and Yamashita, K. 2004. β1,3-N-Acetylglucosaminyltransferase-7 (β3Gn-T7) acts efficiently on keratan sulfate-related glycans. *FEBS Lett.* 556: 216-220.
3. Iwai, T., Kudo, T., Kawamoto, R., Kubota, T., Togayachi, A., Hiruma, T., Okada, T., Kawamoto, T., Morozumi, K. and Narimatsu, H. 2005. Core 3 synthase is down-regulated in colon carcinoma and profoundly suppresses the metastatic potential of carcinoma cells. *Proc. Natl. Acad. Sci. USA* 102: 4572-4577.
4. Deo, V.K. and Park, E.Y. 2006. Multiple co-transfection and co-expression of human β-1,3-N-acetylglucosaminyltransferase with human calreticulin chaperone cDNA in a single step in insect cells. *Biotechnol. Appl. Biochem.* 43: 129-135.
5. Seko, A. and Yamashita, K. 2008. Activation of β1,3-N-acetylglucosaminyltransferase-2 (β3Gn-T2) by β3Gn-T8: Possible involvement of β3Gn-T8 in increasing poly-N-acetyllactosamine chains in differentiated HL-60 cells. *J. Biol. Chem.* 283: 33094-33100.
6. Marcos, N.T., Magalhães, A., Ferreira, B., Oliveira, M.J., Carvalho, A.S., Mendes, N., Gilmartin, T., Head, S.R., Figueiredo, C., David, L., Santos-Silva, F. and Reis, C.A. 2008. *Helicobacter pylori* induces β3GnT5 in human gastric cell lines, modulating expression of the SabA ligand sialyl-Lewis x. *J. Clin. Invest.* 118: 2325-2336.

CHROMOSOMAL LOCATION

Genetic locus: B3gnt9 (mouse) mapping to 8 D3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

β3Gn-T9 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see β3Gn-T9 shRNA Plasmid (m): sc-141927-SH and β3Gn-T9 shRNA (m) Lentiviral Particles: sc-141927-V as alternate gene silencing products.

For independent verification of β3Gn-T9 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141927A, sc-141927B and sc-141927C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

β3Gn-T9 siRNA (m) is recommended for the inhibition of β3Gn-T9 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor β3Gn-T9 gene expression knockdown using RT-PCR Primer: β3Gn-T9 (m)-PR: sc-141927-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.