



## cadherin-22 siRNA (m): sc-141973

### BACKGROUND

The cadherins are a family of  $\text{Ca}^{2+}$ -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of structure and morphogenesis. Cadherins each contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. Cadherin-22, also known as CDH22 or PB-cadherin (pituitary and brain cadherin), is an 828 amino acid single-pass type I membrane protein that, characteristic of cadherin proteins, contains five cadherin domains. Expressed predominately in brain, cadherin-22 functions as a  $\text{Ca}^{2+}$ -dependent cell adhesion protein that is thought to play an important role in tissue formation and morphogenesis, specifically in neural cells during the development and maintenance of brain tissue.

### REFERENCES

1. Sugimoto, K., et al. 1996. Molecular cloning and characterization of a newly identified member of the cadherin family, PB-cadherin. *J. Biol. Chem.* 271: 11548-11556.
2. Kremmidiotis, G., et al. 1998. Localization of human cadherin genes to chromosome regions exhibiting cancer-related loss of heterozygosity. *Genomics* 49: 467-471.
3. Kitajima, K., et al. 1999. Expression of a novel type of classic cadherin, PB-cadherin in developing brain and limb buds. *Dev. Dyn.* 215: 206-214.
4. Wu, J., et al. 2003. Expression of a novel factor, short-type PB-cadherin, in Sertoli cells and spermatogenic stem cells of the neonatal rat testis. *J. Endocrinol.* 176: 381-391.
5. Wu, J., et al. 2005. Short-type PB-cadherin promotes survival of gonocytes and activates JAK-STAT signalling. *Dev. Biol.* 284: 437-450.
6. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 609920. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Bento, J.L., et al. 2008. Heterogeneity in gene loci associated with type 2 diabetes on human chromosome 20q13.1. *Genomics* 92: 226-234.

### CHROMOSOMAL LOCATION

Genetic locus: Cdh22 (mouse) mapping to 2 H3.

### PRODUCT

cadherin-22 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see cadherin-22 shRNA Plasmid (m): sc-141973-SH and cadherin-22 shRNA (m) Lentiviral Particles: sc-141973-V as alternate gene silencing products.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}\text{C}$ , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

cadherin-22 siRNA (m) is recommended for the inhibition of cadherin-22 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor cadherin-22 gene expression knockdown using RT-PCR Primer: cadherin-22 (m)-PR: sc-141973-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^{\circ}\text{C}$  and the extension temperature should be  $68-72^{\circ}\text{C}$ .

### RESEARCH USE

For research use only, not for use in diagnostic procedures.