

COL4A4 siRNA (m): sc-142468

BACKGROUND

The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial collagens (types I, II, III and V) and basement membrane collagens (type IV), each type containing multiple isoforms. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function. COL4A4 (collagen α -4(IV) chain), also known as CA44, is a 1,690 amino acid extracellular matrix protein that functions in the structure of glomerular basement membranes. A member of the type IV collagen family, COL4A4 contains one collagen IV NC1 (C-terminal non-collagenous) domain and is encoded by a gene that maps to human chromosome 2q36.3.

REFERENCES

1. Kamagata, Y., Mattei, M.G. and Ninomiya, Y. 1992. Isolation and sequencing of cDNAs and genomic DNAs encoding the α 4 chain of basement membrane collagen type IV and assignment of the gene to the distal long arm of human chromosome 2. *J. Biol. Chem.* 267: 23753-23758.
2. Mariyama, M., Leinonen, A., Mochizuki, T., Tryggvason, K. and Reenders, S.T. 1994. Complete primary structure of the human α 3(IV) collagen chain. Coexpression of the α 3(IV) and α 4(IV) collagen chains in human tissues. *J. Biol. Chem.* 269: 23013-23017.
3. Leinonen, A., Mariyama, M., Mochizuki, T., Tryggvason, K. and Reenders, S.T. 1994. Complete primary structure of the human type IV collagen α 4(IV) chain. Comparison with structure and expression of the other α (IV) chains. *J. Biol. Chem.* 269: 26172-26177.
4. Lemmink, H.H., Nillesen, W.N., Mochizuki, T., Schröder, C.H., Brunner, H.G., van Oost, B.A., Monnens, L.A. and Smeets, H.J. 1996. Benign familial hematuria due to mutation of the type IV collagen α 4 gene. *J. Clin. Invest.* 98: 1114-1118.
5. Boye, E., Mollet, G., Forestier, L., Cohen-Solal, L., Heidet, L., Cochat, P., Grünfeld, J.P., Palcoux, J.B., Gubler, M.C. and Antignac, C. 1998. Determination of the genomic structure of the COL4A4 gene and of novel mutations causing autosomal recessive Alport syndrome. *Am. J. Hum. Genet.* 63: 1329-1340.
6. Momota, R., Sugimoto, M., Ohashi, T., Kigasawa, K., Yoshioka, H. and Ninomiya, Y. 1998. Two genes, COL4A3 and COL4A4 coding for the human α 3(IV) and α 4(IV) collagen chains are arranged head-to-head on chromosome 2q36. *FEBS Lett.* 424: 11-16.
7. Badenas, C., Praga, M., Tazón, B., Heidet, L., Arrondel, C., Armengol, A., Andres, A., Morales, E., Camacho, J.A., Lens, X., Dávila, S., Milà, M., Antignac, C., Darnell, A. and Torra, R. 2002. Mutations in the COL4A4 and COL4A3 genes cause familial benign hematuria. *J. Am. Soc. Nephrol.* 13: 1248-1254.

CHROMOSOMAL LOCATION

Genetic locus: Col4a4 (mouse) mapping to 1 C5.

PRODUCT

COL4A4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see COL4A4 shRNA Plasmid (m): sc-142468-SH and COL4A4 shRNA (m) Lentiviral Particles: sc-142468-V as alternate gene silencing products.

For independent verification of COL4A4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142468A, sc-142468B and sc-142468C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

COL4A4 siRNA (m) is recommended for the inhibition of COL4A4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor COL4A4 gene expression knockdown using RT-PCR Primer: COL4A4 (m)-PR: sc-142468-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.