

# Cyclophilin G siRNA (m): sc-142661

## BACKGROUND

Cyclophilins are conserved, ubiquitous and abundant cytosolic peptidyl-prolyl *cis-trans* isomerases that accelerate the isomerization of XaaPro peptide bonds and the refolding of proteins. Cyclophilin G, also known as CARS-Cyp (clk-associating RS-cyclophilin), SRcyp (SR-cyclophilin), CASP10, PPlase G or Rotamase G, is a ubiquitously expressed member of the Moca family of cyclophilins. Localizing to the nucleus and, during interphase, nuclear speckles, Cyclophilin G contains an N-terminal cyclophilin-type domain, an acidic serine-rich region, five Cdc2-type phosphorylation sites and a series of serine-arginine repeats throughout the C-terminus. Cyclophilin G is phosphorylated during mitosis by the Cdc2-cyclin B complex, suggesting that its function is cell cycle-regulated. In addition, Cyclophilin G is capable of interacting with Pinin and the C-terminus of the largest subunit of RNA polymerase II (Pol II). Cyclophilin G may participate in pre-mRNA splicing by regulating the subnuclear localization of SR/SR-like protein family members.

## REFERENCES

1. Nestel, F.P., et al. 1996. RS cyclophilins: identification of an NK-TR1-related cyclophilin. *Gene* 180: 151-155.
2. Giardina, S.L., et al. 1996. Association of the expression of an SR-cyclophilin with myeloid cell differentiation. *Blood* 87: 2269-2274.
3. Bourquin, J.P., et al. 1997. A serine/arginine-rich nuclear matrix cyclophilin interacts with the C-terminal domain of RNA polymerase II. *Nucleic Acids Res.* 25: 2055-2061.
4. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606093. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Lin, C.L., et al. 2004. Over-expression of SR-cyclophilin, an interaction partner of nuclear pinin, releases SR family splicing factors from nuclear speckles. *Biochem. Biophys. Res. Commun.* 321: 638-647.

## CHROMOSOMAL LOCATION

Genetic locus: Pp1g (mouse) mapping to 2 C2.

## PRODUCT

Cyclophilin G siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Cyclophilin G shRNA Plasmid (m): sc-142661-SH and Cyclophilin G shRNA (m) Lentiviral Particles: sc-142661-V as alternate gene silencing products.

For independent verification of Cyclophilin G (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142661A, sc-142661B and sc-142661C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Cyclophilin G siRNA (m) is recommended for the inhibition of Cyclophilin G expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Cyclophilin G (8S5): sc-100699 is recommended as a control antibody for monitoring of Cyclophilin G gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Cyclophilin G gene expression knockdown using RT-PCR Primer: Cyclophilin G (m)-PR: sc-142661-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.