

CYP4A10 siRNA (m): sc-142720

BACKGROUND

Cytochrome P450 proteins are heme-thiolate monooxygenases that mediate NADPH-dependent electron transport and function to oxidize a variety of structurally unrelated compounds, including steroids, fatty acids and xenobiotics. Specifically, Cytochrome P450s are responsible for metabolizing arachidonic acid to hydroxyeicosatetraenoic acid (a regulator of blood pressure) and epoxyeicosatrienoic acid (a molecule involved in signaling events). CYP4A10 (cytochrome P450, family 4, subfamily a, polypeptide 10), also known as RP1, cytochrome P452, lauric acid omega-hydroxylase, CYP1A10, Cyp4a or D4Rp1, is a 509 amino acid peripheral membrane protein of the Endoplasmic reticulum and microsome that is encoded by a gene that maps to murine chromosome 4 D1. CYP4A10 is the mouse homolog of human CYP4A11.

REFERENCES

1. Imaoka, S., et al. 1993. Complete cDNA sequence and cDNA-directed expression of CYP4A11, a fatty acid ω -hydroxylase expressed in human kidney. *DNA Cell Biol.* 12: 893-899.
2. Powell, P.K., et al. 1998. Metabolism of arachidonic acid to 20-hydroxy-5,8,11, 14-eicosatetraenoic acid by P450 enzymes in human liver: involvement of CYP4F2 and CYP4A11. *J. Pharmacol. Exp. Ther.* 285: 1327-1336.
3. Chang, Y.T., et al. 1999. Homology modeling and substrate binding study of human CYP4A11 enzyme. *Proteins* 34: 403-415.
4. Lasker, J.M., et al. 2000. Formation of 20-hydroxyeicosatetraenoic acid, a vasoactive and natriuretic eicosanoid, in human kidney. Role of Cyp4F2 and Cyp4A11. *J. Biol. Chem.* 275: 4118-4126.
5. Hoch, U., et al. 2001. Covalently linked heme in cytochrome p4504a fatty acid hydroxylases. *J. Biol. Chem.* 276: 11339-11346.
6. Gainer, J.V., et al. 2005. Functional variant of CYP4A11 20-hydroxyeicosatetraenoic acid synthase is associated with essential hypertension. *Circulation* 111: 63-69.
7. Fu, Z., et al. 2008. Haplotype-based case study of human CYP4A11 gene and cerebral infarction in Japanese subject. *Endocrine* 33: 215-222.

CHROMOSOMAL LOCATION

Genetic locus: Cyp4a10 (mouse) mapping to 4 D1.

PRODUCT

CYP4A10 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CYP4A10 shRNA Plasmid (m): sc-142720-SH and CYP4A10 shRNA (m) Lentiviral Particles: sc-142720-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CYP4A10 siRNA (m) is recommended for the inhibition of CYP4A10 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CYP4A10 gene expression knockdown using RT-PCR Primer: CYP4A10 (m)-PR: sc-142720-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.