

# DCAMKL3 siRNA (m): sc-142890

## BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including cell division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine (Ser/Thr) protein kinases. DCAMKL3 (doublecortin-like and CAM kinase-like 3), also known as DCLK3 or DCDC3C, is a 648 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one protein kinase domain. One of several members of the Ser/Thr protein kinase family, DCAMKL3 functions to catalyze the ATP-dependent attachment of a phospho residue to target proteins, an event that may play a role in various cellular processes.

## REFERENCES

1. Bairoch, A., et al. 1988. Sequence patterns in protein kinases. *Nature* 331: 22.
2. Hanks, S.K., et al. 1988. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241: 42-52.
3. Hanks, S.K., et al. 1991. Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. *Meth. Enzymol.* 200: 38-62.
4. Sapir, T., et al. 2000. Doublecortin mutations cluster in evolutionarily conserved functional domains. *Hum. Mol. Genet.* 9: 703-712.
5. Reiner, O., et al. 2006. The evolving doublecortin (DCX) superfamily. *BMC Genomics* 7: 188.
6. Coquelle, F.M., et al. 2006. Common and divergent roles for members of the mouse DCX superfamily. *Cell Cycle* 5: 976-983.
7. Cierpicki, T., et al. 2006. The DC-module of doublecortin: dynamics, domain boundaries, and functional implications. *Proteins* 64: 874-882.

## CHROMOSOMAL LOCATION

Genetic locus: Dclk3 (mouse) mapping to 9 F3.

## PRODUCT

DCAMKL3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DCAMKL3 shRNA Plasmid (m): sc-142890-SH and DCAMKL3 shRNA (m) Lentiviral Particles: sc-142890-V as alternate gene silencing products.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

DCAMKL3 siRNA (m) is recommended for the inhibition of DCAMKL3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DCAMKL3 gene expression knockdown using RT-PCR Primer: DCAMKL3 (m)-PR: sc-142890-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.