

DDX35 siRNA (m): sc-142932

BACKGROUND

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp, are putative RNA helicases implicated in several cellular processes involving modifications of RNA secondary structure. Specifically, DEAD box proteins are involved in translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, members of this family may be involved in embryogenesis, spermatogenesis, and cellular growth and division. DDX30, DDX35 and DDX36 each contain 1 helicase ATP-binding domain and 1 helicase C-terminal domain. DDX35 (DEAD box protein 35) is a 703 amino acid protein that is part of the spliceosome C complex and is involved in pre-mRNA splicing.

REFERENCES

1. Fu, J.J., et al. 2002. Molecular cloning and characterization of human DDX36 and mouse Ddx36 genes, new members of the DEAD/H box superfamily. *Sheng Wu Hua Xue Yu Sheng Wu Wu Li Xue Bao* 34: 655-661.
2. Jurica, M.S., et al. 2002. Purification and characterization of native spliceosomes suitable for three-dimensional structural analysis. *RNA* 8: 426-439.
3. Fu, J.J., et al. 2003. Expression research for human DDX36 and mouse Ddx36 gene in the adult testis. *Yi Chuan Xue Bao* 30: 201-208.
4. Abdelhaleem, M. 2005. RNA helicases: regulators of differentiation. *Clin. Biochem.* 38: 499-503.
5. Cordin, O., et al. 2006. The DEAD-box protein family of RNA helicases. *Gene* 367: 17-37.
6. Wang, Y., et al. 2006. Human mitochondrial DNA nucleoids are linked to protein folding machinery and metabolic enzymes at the mitochondrial inner membrane. *J. Biol. Chem.* 281: 25791-25802.

CHROMOSOMAL LOCATION

Genetic locus: Dhx35 (mouse) mapping to 2 H1.

PRODUCT

DDX35 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DDX35 shRNA Plasmid (m): sc-142932-SH and DDX35 shRNA (m) Lentiviral Particles: sc-142932-V as alternate gene silencing products.

For independent verification of DDX35 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142932A, sc-142932B and sc-142932C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DDX35 siRNA (m) is recommended for the inhibition of DDX35 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DDX35 gene expression knockdown using RT-PCR Primer: DDX35 (m)-PR: sc-142932-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.