

DnaJC4 siRNA (m): sc-143109

BACKGROUND

The DnaJ family is one of the largest of all chaperone families and has evolved with diverse cellular localization and functions. Presence of a J domain defines a protein as a member of the DnaJ family. DnaJ heat shock induced proteins are derived from *Escherichia coli* and are under the control of the htpR regulatory protein. DnaJ proteins play a critical role in the HSP 70 chaperone machine by interacting with HSP 70 to stimulate ATP hydrolysis. DnaJ proteins contain cysteine rich regions that are composed of zinc fingers, which form a peptide binding domain responsible for the chaperone function. DnaJ proteins are important mediators of proteolysis and are involved in the regulation of protein degradation, exocytosis and endocytosis. DnaJC4 (DnaJ homolog subfamily C member 4), also known as HSPF2, MCG18 or DANJC4, is a 241 amino acid membrane protein that contains one J domain.

REFERENCES

1. Saito, H., et al. 1978. Organization and expression of the DnaJ and DnaK genes of *Escherichia coli* K12. Mol. Gen. Genet. 164: 1-8.
2. Georgopoulos, C.P., et al. 1980. Identification of the *E. coli* dnaJ gene product. Mol. Gen. Genet. 178: 583-588.
3. Suh, W.C., et al. 1998. Interaction of the Hsp70 molecular chaperone, DnaK, with its cochaperone DnaJ. Proc. Natl. Acad. Sci. USA 95: 15223-15228.
4. Tomoyasu, T., et al. 1998. Levels of DnaK and DnaJ provide tight control of heat shock gene expression and protein repair in *Escherichia coli*. Mol. Microbiol. 30: 567-581.
5. Stewart, G.R., et al. 2004. Analysis of the function of mycobacterial DnaJ proteins by overexpression and microarray profiling. Tuberculosis 84: 180-187.
6. Shi, Y.Y., et al. 2005. The C-terminal (331-376) sequence of *Escherichia coli* DnaJ is essential for dimerization and chaperone activity: a small angle X-ray scattering study in solution. J. Biol. Chem. 280: 22761-22768.
7. Robichon, C., et al. 2006. DnaJA4 is a SREBP-regulated chaperone involved in the cholesterol biosynthesis pathway. Biochim. Biophys. Acta 1761: 1107-1113.
8. Acebrón, S.P., et al. 2008. DnaJ recruits DnaK to protein aggregates. J. Biol. Chem. 283: 1381-1390.

CHROMOSOMAL LOCATION

Genetic locus: Dnajc4 (mouse) mapping to 19 A.

PRODUCT

DnaJC4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DnaJC4 shRNA Plasmid (m): sc-143109-SH and DnaJC4 shRNA (m) Lentiviral Particles: sc-143109-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DnaJC4 siRNA (m) is recommended for the inhibition of DnaJC4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DnaJC4 gene expression knockdown using RT-PCR Primer: DnaJC4 (m)-PR: sc-143109-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.