

DNAL4 siRNA (m): sc-143115

BACKGROUND

Dyneins are multisubunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Belonging to the Dynein light chain family, DNAL4 (dynein light chain 4, axonemal) is a 105 amino acid protein that is a component of the Dynein motor complex. Cytoplasmic or axonemal Dynein heavy, intermediate, light and light-intermediate chains are all components of minus end-directed motors; complexes that transport cellular cargos towards the central region of the cell. Axonemal Dynein motors contain one to three non-identical heavy chains and cause a sliding of microtubules in the axonemes of cilia and flagella in a mechanism necessary for cilia to beat and propel the cell.

REFERENCES

1. Vaughan, K.T., Mikami, A., Paschal, B.M., Holzbaur, E.L., Hughes, S.M., Echeverri, C.J., Moore, K.J., Gilbert, D.J., Copeland, N.G., Jenkins, N.A. and Vallee, R.B. 1996. Multiple mouse chromosomal loci for dynein-based motility. *Genomics* 36: 29-38.
2. Yagi, T. 2000. ADP-dependent microtubule translocation by flagellar inner-arm dyneins. *Cell Struct. Funct.* 25: 263-267.
3. Epstein, E., Sela-Brown, A., Ringel, I., Kilav, R., King, S.M., Benashski, S.E., Yisraeli, J.K., Silver, J. and Naveh-Manny, T. 2000. Dynein light chain binding to a 3'-untranslated sequence mediates parathyroid hormone mRNA association with microtubules. *J. Clin. Invest.* 105: 505-512.
4. Tang, Q., Staub, C.M., Gao, G., Jin, Q., Wang, Z., Ding, W., Aurigemma, R.E. and Mulder, K.M. 2002. A novel transforming growth factor- β receptor-interacting protein that is also a light chain of the motor protein dynein. *Mol. Biol. Cell* 13: 4484-4496.
5. Wu, H. and King, S.M. 2003. Backbone dynamics of dynein light chains. *Cell Motil. Cytoskeleton* 54: 267-273.
6. Pazour, G.J., Agrin, N., Walker, B.L. and Witman, G.B. 2006. Identification of predicted human outer dynein arm genes: candidates for primary ciliary dyskinesia genes. *J. Med. Genet.* 43: 62-73.
7. Benison, G., Karplus, P.A. and Barbar, E. 2008. The interplay of ligand binding and quaternary structure in the diverse interactions of dynein light chain LC8. *J. Mol. Biol.* 384: 954-966.
8. Tanner, C.A., Rompolas, P., Patel-King, R.S., Gorbatyuk, O., Wakabayashi, K., Pazour, G.J. and King, S.M. 2008. Three members of the LC8/DYNLL family are required for outer arm dynein motor function. *Mol. Biol. Cell* 19: 3724-3734.

CHROMOSOMAL LOCATION

Genetic locus: Dnal4 (mouse) mapping to 15 E1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

DNAL4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DNAL4 shRNA Plasmid (m): sc-143115-SH and DNAL4 shRNA (m) Lentiviral Particles: sc-143115-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DNAL4 siRNA (m) is recommended for the inhibition of DNAL4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DNAL4 gene expression knockdown using RT-PCR Primer: DNAL4 (m)-PR: sc-143115-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.