

DNTTIP2 siRNA (m): sc-143130

BACKGROUND

DNTTIP2 (deoxynucleotidyltransferase terminal-interacting protein 2), also known as ERBP (estrogen receptor-binding protein), FCF2 or TdIF2 (terminal deoxynucleotidyltransferase-interacting factor 2), is a 756 amino acid nuclear protein. Widely expressed, with highest levels in testis, DNTTIP2 controls the transcriptional activity of TdT and ER α . DNTTIP2 forms a ternary complex with TdT and core histones. DNTTIP2 may also function as a chromatin remodeling protein. DNTTIP2 is encoded by a gene that maps to human chromosome 1p22.1, which is the largest human chromosome, spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes Lamin A. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

1. Watson, M.L., Kingsmore, S.F., Johnston, G.I., Siegelman, M.H., Le Beau, M.M., Lemons, R.S., Bora, N.S., Howard, T.A., Weissman, I.L., McEver, R.P., et al. 1990. Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263-272.
2. Soares-da-Silva, P. and Fernandes, M.H. 1992. Sodium-dependence and ouabain-sensitivity of the synthesis of dopamine in renal tissues of the rat. *Br. J. Pharmacol.* 105: 811-816.
3. Blackwood, D.H., Fordyce, A., Walker, M.T., St Clair, D.M., Porteous, D.J. and Muir, W.J. 2001. Schizophrenia and affective disorders—cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am. J. Hum. Genet.* 69: 428-433.
4. Fujita, K., Shimazaki, N., Ohta, Y., Kubota, T., Ibe, S., Toji, S., Tamai, K., Fujisaki, S., Hayano, T. and Koiwai, O. 2003. Terminal deoxynucleotidyltransferase forms a ternary complex with a novel chromatin remodeling protein with 82 kDa and core histone. *Genes Cells* 8: 559-571.
5. Bu, H., Kashireddy, P., Chang, J., Zhu, Y.T., Zhang, Z., Zheng, W., Rao, S.M. and Zhu, Y.J. 2004. ERBP, a novel estrogen receptor binding protein enhancing the activity of estrogen receptor. *Biochem. Biophys. Res. Commun.* 317: 54-59.

CHROMOSOMAL LOCATION

Genetic locus: Dnttip2 (mouse) mapping to 3 G1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

DNTTIP2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DNTTIP2 shRNA Plasmid (m): sc-143130-SH and DNTTIP2 shRNA (m) Lentiviral Particles: sc-143130-V as alternate gene silencing products.

For independent verification of DNTTIP2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-143130A, sc-143130B and sc-143130C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DNTTIP2 siRNA (m) is recommended for the inhibition of DNTTIP2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DNTTIP2 gene expression knockdown using RT-PCR Primer: DNTTIP2 (m)-PR: sc-143130-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.