

## FAT4 siRNA (m): sc-145082

### BACKGROUND

The cadherins represent a family of  $\text{Ca}^{2+}$ -dependent adhesion molecules that function to mediate cell to cell binding that is critical for the maintenance of structure and morphogenesis. Cadherins contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short C-terminal intracellular domain interacts with a variety of cytoplasmic proteins, including  $\beta$ -catenin, to regulate cadherin function. The cadherin superfamily includes cadherins, protocadherins, desmogleins and desmocollins. FAT4 (FAT tumor suppressor homolog 4), also known as FAT-J, CDHF14 or CDHR11, is a 4,981 amino acid single-pass type I membrane protein that belongs to the protocadherin subfamily of cadherins and localizes to the primary cilia of kidney. Widely expressed, FAT4 contains thirty-four cadherin domains, six EGF-like domains and two laminin G-like domains. FAT4 may function in the regulation of planar cell polarity.

### REFERENCES

1. Sano, K., Tanihara, H., Heimark, R.L., Obata, S., Davidson, M., St. John, T., Taketani, S. and Suzuki, S. 1993. Protocadherins: a large family of cadherin-related molecules in central nervous system. *EMBO J.* 12: 2249-2256.
2. Wu, Q. and Maniatis, T. 1999. A striking organization of a large family of human neural cadherin-like cell adhesion genes. *Cell* 97: 779-790.
3. Suzuki, S.T. 2000. Recent progress in protocadherin research. *Exp. Cell Res.* 261: 13-18.
4. Hill, E., Broadbent, I.D., Chothia, C. and Pettitt, J. 2001. Cadherin superfamily proteins in *Caenorhabditis elegans* and *Drosophila melanogaster*. *J. Mol. Biol.* 305: 1011-1024.
5. Wolverton, T. and Lalande, M. 2001. Identification and characterization of three members of a novel subclass of protocadherins. *Genomics* 76: 66-72.
6. Noonan, J.P., Grimwood, J., Schmutz, J., Dickson, M. and Myers, R.M. 2004. Gene conversion and the evolution of protocadherin gene cluster diversity. *Genome Res.* 14: 354-366.
7. Durand, C.M., Kappeler, C., Betancur, C., Delorme, R., Quach, H., Goubran-Botros, H., Melke, J., Nygren, G., Chabane, N., Bellivier, F., Szoke, A., Schurhoff, F., Rastam, M., Anckarsäter, H., Gillberg, C., Leboyer, M. et al. 2006. Expression and genetic variability of PCDH11Y, a gene specific to Homo sapiens and candidate for susceptibility to psychiatric disorders. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 141: 67-70.

### CHROMOSOMAL LOCATION

Genetic locus: Fat4 (mouse) mapping to 3 B.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

### PRODUCT

FAT4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FAT4 shRNA Plasmid (m): sc-145082-SH and FAT4 shRNA (m) Lentiviral Particles: sc-145082-V as alternate gene silencing products.

For independent verification of FAT4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145082A, sc-145082B and sc-145082C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}\text{C}$ , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

FAT4 siRNA (m) is recommended for the inhibition of FAT4 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FAT4 gene expression knockdown using RT-PCR Primer: FAT4 (m)-PR: sc-145082-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^{\circ}\text{C}$  and the extension temperature should be  $68-72^{\circ}\text{C}$ .