

FBXO27 siRNA (m): sc-145113

BACKGROUND

In eukaryotes, degradation of damaged or excess proteins into short peptides is carried out by proteasomes. The proteasomes bind polyubiquitin chains that are added to the target proteins through a phosphorylation-dependent reaction catalyzed by ubiquitin ligases, such as the SCF-type E3 complex containing Skp, Cullin, Rbx1 and F-box proteins. F-box proteins, such as FBXO27 (F-box only protein 27), possess structural motifs used for directly aggregating the substrate while binding to the Skp1 bridge providing for close proximity to the functional E2 ubiquitin-conjugating enzyme, Cullin/Rbx1. FBXO27, also known as FBG5 and FBX27, is a 283 amino acid protein that contains an N-terminal F-box and a C-terminal F-box-associated domain. FBXO27 found highly expressed in brain, heart and kidney with lower levels found liver.

REFERENCES

1. Cenciarelli, C., Chiaur, D.S., Guardavaccaro, D., Parks, W., Vidal, M. and Pagano, M. 1999. Identification of a family of human F-box proteins. *Curr. Biol.* 9: 1177-1179.
2. Winston, J.T., Koepp, D.M., Zhu, C., Elledge, S.J. and Harper, J.W. 1999. A family of mammalian F-box proteins. *Curr. Biol.* 9: 1180-1182.
3. Craig, K.L. and Tyers, M. 1999. The F-box: a new motif for ubiquitin dependent proteolysis in cell cycle regulation and signal transduction. *Prog. Biophys. Mol. Biol.* 72: 299-328.
4. Ilyin, G.P., Rialland, M., Pigeon, C. and Guguen-Guillouzo, C. 2000. cDNA cloning and expression analysis of new members of the mammalian F-box protein family. *Genomics* 67: 40-47.
5. Ilyin, G.P., Serandour, A.L., Pigeon, C., Rialland, M., Glaise, D. and Guguen-Guillouzo, C. 2002. A new subfamily of structurally related human F-box proteins. *Gene* 296: 11-20.
6. Glenn, K.A., Nelson, R.F., Wen, H.M., Mallinger, A.J. and Paulson, H.L. 2008. Diversity in tissue expression, substrate binding, and SCF complex formation for a lectin family of ubiquitin ligases. *J. Biol. Chem.* 283: 12717-12729.

CHROMOSOMAL LOCATION

Genetic locus: Fbxo27 (mouse) mapping to 7 A3.

PRODUCT

FBXO27 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FBXO27 shRNA Plasmid (m): sc-145113-SH and FBXO27 shRNA (m) Lentiviral Particles: sc-145113-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FBXO27 siRNA (m) is recommended for the inhibition of FBXO27 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FBXO27 gene expression knockdown using RT-PCR Primer: FBXO27 (m)-PR: sc-145113-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.