

FBXW5 siRNA (m): sc-145143

BACKGROUND

F-box proteins are critical components of the SCF (Skp1-CUL-1-F-box protein) type E3 ubiquitin ligase complex and are involved in substrate recognition and recruitment for ubiquitination. They are members of a larger family of proteins that are involved in the regulation of a wide variety of cellular processes (including the cell cycle, immune responses, signaling cascades and developmental events) through the targeting of proteins, such as cyclins, cyclin-dependent kinase inhibitors, I κ B- α and β -catenin, for proteasomal degradation. FBXW5 (F-box and WD repeat domain containing 5), also known as FBW5, is a 566 amino acid protein that contains one F-box domain and three WD repeats. Expressed as multiple alternatively spliced isoforms, FBXW5 functions as a substrate-recognition component of the SCF complex and directly interacts with Skp1 p19 and CUL-1.

REFERENCES

1. Bai, C., Sen, P., Hofmann, K., Ma, L., Goebel, M., Harper, J.W. and Elledge, S.J. 1996. Skp1 connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell* 86: 263-274.
2. Skowyra, D., Craig, K.L., Tyers, M., Elledge, S.J. and Harper, J.W. 1997. F-box proteins are receptors that recruit phosphorylated substrates to the SCF ubiquitin-ligase complex. *Cell* 91: 209-219.
3. Winston, J.T., Koepp, D.M., Zhu, C., Elledge, S.J. and Harper, J.W. 1999. A family of mammalian F-box proteins. *Curr. Biol.* 9: 1180-1182.
4. Craig, K.L. and Tyers, M. 1999. The F-box: a new motif for ubiquitin dependent proteolysis in cell cycle regulation and signal transduction. *Prog. Biophys. Mol. Biol.* 72: 299-328.
5. Kipreos, E.T. and Pagano, M. 2000. The F-box protein family. *Genome Biol.* 1: S3002.
6. Ilyin, G.P., Serandour, A.L., Pigeon, C., Rialland, M., Glaise, D. and Guguen-Guillouzo, C. 2002. A new subfamily of structurally related human F-box proteins. *Gene* 296: 11-20.
7. Hu, J., Zacharek, S., He, Y.J., Lee, H., Shumway, S., Duronio, R.J. and Xiong, Y. 2008. WD40 protein FBW5 promotes ubiquitination of tumor suppressor TSC2 by DDB1-CUL-4-ROC1 ligase. *Genes Dev.* 22: 866-871.
8. Glenn, K.A., Nelson, R.F., Wen, H.M., Mallinger, A.J. and Paulson, H.L. 2008. Diversity in tissue expression, substrate binding, and SCF complex formation for a lectin family of ubiquitin ligases. *J. Biol. Chem.* 283: 12717-12729.
9. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 609072. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Fbxw5 (mouse) mapping to 2 A3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

FBXW5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FBXW5 shRNA Plasmid (m): sc-145143-SH and FBXW5 shRNA (m) Lentiviral Particles: sc-145143-V as alternate gene silencing products.

For independent verification of FBXW5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145143A, sc-145143B and sc-145143C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FBXW5 siRNA (m) is recommended for the inhibition of FBXW5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FBXW5 gene expression knockdown using RT-PCR Primer: FBXW5 (m)-PR: sc-145143-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.