

FCAMR siRNA (m): sc-145146

BACKGROUND

Fc (Ig constant fragment) receptors ensure protection of the host against foreign antigens, such as microorganisms and pathogens, by removing Ig-coated antigen complexes from circulation. Fc receptors are present on lymphoid and myeloid derivatives, where they mediate endocytosis of Ig-antigen complexes, antibody production in B cells through T cell antigen presentation, cytotoxicity and the release of cytokines and reactive oxygen species. The Fc α/μ receptor (FCAMR) functions as a receptor for the Fc fragment of IgA and IgM and mediates their endocytosis. FCAMR is a 532 amino acid protein that is expressed in mesangial cells and is upregulated by IFNG/IFN- γ . It is also expressed in secondary lymphoid organs, such as lymph node and appendix, kidney and intestine, suggesting an important role of the receptor for immunity in these organs.

REFERENCES

- Shibuya, A., Sakamoto, N., Shimizu, Y., Shibuya, K., Osawa, M., Hiroyama, T., Eyre, H.J., Sutherland, G.R., Endo, Y., Fujita, T., Miyabayashi, T., Sakano, S., Tsuji, T., Nakayama, E., Phillips, J.H., Lanier, L.L. and Nakauchi, H. 2000. Fc α/μ receptor mediates endocytosis of IgM-coated microbes. *Nat. Immunol.* 1: 441-446.
- Sakamoto, N., Shibuya, K., Shimizu, Y., Yotsumoto, K., Miyabayashi, T., Sakano, S., Tsuji, T., Nakayama, E., Nakauchi, H. and Shibuya, A. 2001. A novel Fc receptor for IgA and IgM is expressed on both hematopoietic and non-hematopoietic tissues. *Eur. J. Immunol.* 31: 1310-1316.
- Shimizu, Y., Honda, S., Yotsumoto, K., Tahara-Hanaoka, S., Eyre, H.J., Sutherland, G.R., Endo, Y., Shibuya, K., Koyama, A., Nakauchi, H. and Shibuya, A. 2001. Fc(α/μ) receptor is a single gene-family member closely related to polymeric immunoglobulin receptor encoded on Chromosome 1. *Immunogenetics* 53: 709-711.
- McDonald, K.J., Cameron, A.J., Allen, J.M. and Jardine, A.G. 2002. Expression of Fc α/μ receptor by human mesangial cells: a candidate receptor for immune complex deposition in IgA nephropathy. *Biochem. Biophys. Res. Commun.* 290: 438-442.
- Hu, R.H., Zhang, Y. and Zhao, M.H. 2005. Transferrin receptor and Fc α/μ receptor may not be the major IgA₁ receptor on human mesangial cells. *Chin. Med. J.* 118: 781-785.
- Brown, E.E., Edberg, J.C. and Kimberly, R.P. 2007. Fc receptor genes and the systemic lupus erythematosus diathesis. *Autoimmunity* 40: 567-581.

CHROMOSOMAL LOCATION

Genetic locus: Fcarm (mouse) mapping to 1 E4.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

FCAMR siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FCAMR shRNA Plasmid (m): sc-145146-SH and FCAMR shRNA (m) Lentiviral Particles: sc-145146-V as alternate gene silencing products.

For independent verification of FCAMR (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145146A, sc-145146B and sc-145146C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FCAMR siRNA (m) is recommended for the inhibition of FCAMR expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FCAMR gene expression knockdown using RT-PCR Primer: FCAMR (m)-PR: sc-145146-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.