

G_γ 11 siRNA (m): sc-145284

BACKGROUND

Members of the guanine nucleotide-binding protein (G protein) γ family directly regulate various activities of ion channels and enzymes. Eight known human G protein γ subunits exist, three of which are novel forms that are designated G_γ 4, G_γ 10 and G_γ 11. G_γ 11 (guanine nucleotide binding protein (G protein), γ 11), also known as GNGT11 or GNG11, is a 73 amino acid lipid-anchored, cell membrane protein belonging to the G protein γ family. G_γ 11 is essential for GTPase activity, G protein-effector interaction and replacement of GDP by GTP. Involved in transmembrane signaling and cellular senescence, G_γ 11 is abundantly expressed in most tissues (with the exception of brain). Decreased expression of G_γ 11 may be linked to splenic marginal zone lymphomas, and the gene encoding G_γ 11 maps to human chromosome 7q21.3.

REFERENCES

1. Cali, J.J., Balcueva, E.A., Rybalkin, I. and Robishaw, J.D. 1992. Selective tissue distribution of G protein γ subunits, including a new form of the γ subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.
2. Ray, K., Kunsch, C., Bonner, L.M. and Robishaw, J.D. 1995. Isolation of cDNA clones encoding eight different human G protein γ subunits, including three novel forms designated the γ 4, γ 10 and γ 11 subunits. *J. Biol. Chem.* 270: 21765-21771.
3. Downes, G.B. and Gautam, N. 1999. The G protein subunit gene families. *Genomics* 62: 544-552.
4. Online Mendelian Inheritance in Man, OMIM[™]. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 604390. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Cuello, F., Schulze, R.A., Heemeyer, F., Meyer, H.E., Lutz, S., Jakobs, K.H., Niroomand, F. and Wieland, T. 2003. Activation of heterotrimeric G proteins by a high energy phosphate transfer via nucleoside diphosphate kinase (NDPK) B and G_β subunits. Complex formation of NDPK B with G_β γ dimers and phosphorylation of His-266 IN G_β. *J. Biol. Chem.* 278: 7220-7226.
6. Ruiz-Ballesteros, E., Mollejo, M., Rodriguez, A., Camacho, F.I., Algara, P., Martinez, N., Pollán, M., Sanchez-Aguilera, A., Menarguez, J., Campo, E., Martinez, P., Mateo, M. and Piris, M.A. 2005. Splenic marginal zone lymphoma: proposal of new diagnostic and prognostic markers identified after tissue and cDNA microarray analysis. *Blood* 106: 1831-1838.
7. Hossain, M.N., Sakemura, R., Fujii, M. and Ayusawa, D. 2006. G-protein γ subunit GNG11 strongly regulates cellular senescence. *Biochem. Biophys. Res. Commun.* 351: 645-650.

CHROMOSOMAL LOCATION

Genetic locus: Gng11 (mouse) mapping to 6 A1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

G_γ 11 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see G_γ 11 shRNA Plasmid (m): sc-145284-SH and G_γ 11 shRNA (m) Lentiviral Particles: sc-145284-V as alternate gene silencing products.

For independent verification of G_γ 11 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145284A and sc-145284B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

G_γ 11 siRNA (m) is recommended for the inhibition of G_γ 11 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor G_γ 11 gene expression knockdown using RT-PCR Primer: G_γ 11 (m)-PR: sc-145284-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.