

# GADL1 siRNA (m): sc-145305

## BACKGROUND

There are two forms of glutamic acid decarboxylases (GADs) that exist in brain: GAD-65 (also known as GAD2) and GAD-67 (also known as GAD1, GAD or SCP). GAD-65 and GAD-67 are members of the group II decarboxylase family of proteins and are responsible for catalyzing the rate limiting step in the production of GABA ( $\gamma$ -aminobutyric acid) from L-glutamic acid. Although both GADs are found in brain, GAD-65 localizes to synaptic vesicle membranes in nerve terminals, while GAD-67 is distributed throughout the cell. GAD-67 is responsible for the basal levels of GABA synthesis. In the case of a heightened demand for GABA in neurotransmission, GAD-65 transiently activates to assist in GABA production. As a member of the group II decarboxylase family, GADL1 (Glutamate decarboxylase-like protein 1) is a 521 amino acid protein that utilizes pyridoxal phosphate as a cofactor for its carboxylase activity. There are two isoforms of GADL1 that exist as a result of alternative splicing events.

## REFERENCES

1. Kanter, I.C., Huttner, H.B., Staykov, D., Biermann, T., Struffert, T., Kerling, F., Hilz, M.J., Schellinger, P.D., Schwab, S. and Bardutzky, J. 2007. Cyclophosphamide for anti-GAD antibody-positive refractory status epilepticus. *Epilepsia* 49: 914-920.
2. Korpershoek, E., Verwest, A.M., Ijzendoorn, Y., Rottier, R., Drexhage, H.A. and de Krijger, R.R. 2007. Expression of GAD-67 and novel GAD-67 splice variants during human fetal pancreas development: GAD-67 expression in the fetal pancreas. *Endocr. Pathol.* 18: 31-36.
3. Kanaani, J., Patterson, G., Schaufele, F., Lippincott-Schwartz, J. and Baekkeskov, S. 2008. A palmitoylation cycle dynamically regulates partitioning of the GABA-synthesizing enzyme GAD-65 between ER-Golgi and post-Golgi membranes. *J. Cell Sci.* 121: 437-449.
4. Wei, J. and Wu, J.Y. 2008. Post-translational regulation of L-glutamic acid decarboxylase in the brain. *Neurochem. Res.* 33: 1459-1465.
5. Hwang, I.K., Li, H., Yoo, K.Y., Choi, J.H., Lee, C.H., Chung, D.W., Kim, D.W., Seong, J.K., Yoon, Y.S., Lee, I.S. and Won, M.H. 2008. Comparison of glutamic acid decarboxylase 67 immunoreactive neurons in the hippocampal CA1 region at various age stages in dogs. *Neurosci. Lett.* 431: 251-255.
6. Ito, T., Hioki, H., Nakamura, K., Kaneko, T., Iino, S. and Nojyo, Y. 2008. Some  $\gamma$ -motoneurons contain  $\gamma$ -aminobutyric acid in the rat cervical spinal cord. *Brain Res.* 1201: 78-87.
7. Hamilton, K.A., Parrish-Aungst, S., Margolis, F.L., Erdelyi, F., Szabó, G. and Puche, A.C. 2008. Sensory deafferentation transsynaptically alters neuronal GluR1 expression in the external plexiform layer of the adult mouse main olfactory bulb. *Chem. Senses* 33: 201-210.
8. Jain, R., Tartar, D.M., Gregg, R.K., Divekar, R.D., Bell, J.J., Lee, H.H., Yu, P., Ellis, J.S., Hoeman, C.M., Franklin, C.L. and Zaghoulani, H. 2008. Innocuous IFN $\gamma$  induced by adjuvant-free antigen restores normoglycemia in NOD mice through inhibition of IL-17 production. *J. Exp. Med.* 205: 207-218.

## CHROMOSOMAL LOCATION

Genetic locus: Gad1l (mouse) mapping to 9 F3.

## PRODUCT

GADL1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GADL1 shRNA Plasmid (m): sc-145305-SH and GADL1 shRNA (m) Lentiviral Particles: sc-145305-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

GADL1 siRNA (m) is recommended for the inhibition of GADL1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GADL1 gene expression knockdown using RT-PCR Primer: GADL1 (m)-PR: sc-145305-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.