

GalNAc-TL4 siRNA (m): sc-145315

BACKGROUND

The UDP-N-acetyl- α -D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes are substrate-specific proteins that catalyze the transfer of GalNAc (N-acetylgalactosaminyl) to serine and threonine residues onto various proteins, thereby initiating mucin-type O-linked glycosylation in Golgi apparatus. GalNAc-TL4, also known as LGALS14 or polypeptide GalNAc transferase-like protein 4, is a 607 amino acid single-pass type II membrane protein belonging to the glycosyltransferase 2 family and GalNAc-T subfamily. Localizing to Golgi apparatus, GalNAc-TL4 utilizes manganese and calcium as cofactors and may assist with the transfer of N-acetyl-D-galactosamine to a serine or threonine residue on protein receptors. GalNAc-TL4 likely catalyzes the initial reaction in O-linked oligosaccharide biosynthesis and contains a ricin B-type lectin domain, which binds to GalNAc and contributes to glycopeptide specificity, and two conserved domains located in the glycosyltransferase region. The N-terminal domain, also known as domain A or GT1 motif, may be involved in manganese coordination and substrate binding while the C-terminal domain, also known as domain B or Gal/GalNAc-T motif, is likely involved in catalytic reaction and UDP-Gal binding. GalNAc-TL4 exists as two alternatively spliced isoforms.

REFERENCES

1. Porowska, H., Paszkiewicz-Gadek, A. and Gindziński, A. 1999. Activity of partially purified UDP-N-acetyl- α -D-galactosamine: polypeptide N-acetylgalactosaminyltransferase with different peptide acceptors. *Acta Biochim. Pol.* 46: 365-370.
2. Kumar, S., Connor, J.R., Dodds, R.A., Halsey, W., Van Horn, M., Mao, J., Sathe, G., Mui, P., Agarwal, P., Badger, A.M., Lee, J.C., Gowen, M. and Lark, M.W. 2001. Identification and initial characterization of 5000 expressed sequenced tags (ESTs) each from adult human normal and osteoarthritic cartilage cDNA libraries. *Osteoarthr. Cartil.* 9: 641-653.
3. Schwientek, T., Bennett, E.P., Flores, C., Thacker, J., Hollmann, M., Reis, C.A., Behrens, J., Mandel, U., Keck, B., Schäfer, M.A., Haselmann, K., Zubarev, R., Roepstorff, P., Burchell, J.M., Taylor-Papadimitriou, J., et al. 2002. Functional conservation of subfamilies of putative UDP-N-acetylgalactosamine:polypeptide N-acetylgalactosaminyltransferases in *Drosophila*, *Caenorhabditis elegans*, and mammals. One subfamily composed of I(2)35Aa is essential in *Drosophila*. *J. Biol. Chem.* 277: 22623-22638.
4. Argüeso, P., Tisdale, A., Mandel, U., Letko, E., Foster, C.S. and Gipson, I.K. 2003. The cell-layer- and cell-type-specific distribution of GalNAc-transferases in the ocular surface epithelia is altered during keratinization. *Invest. Ophthalmol. Vis. Sci.* 44: 86-92.
5. Gerhard, D.S., Wagner, L., Feingold, E.A., Shenmen, C.M., Grouse, L.H., Schuler, G., Klein, S.L., Old, S., Rasooly, R., Good, P., Guyer, M., Peck, A.M., Derge, J.G., Lipman, D., Collins, F.S., Jang, W., Sherry, S., et al. 2004. The status, quality, and expansion of the NIH full-length cDNA project: the mammalian gene collection (MGC). *Genome Res.* 14: 2121-2127.
6. Herr, P., Korniyuchuk, G., Yamamoto, Y., Grubisic, K. and Oelgeschläger, M. 2008. Regulation of TGF- β signalling by N-acetylgalactosaminyltransferase-like 1. *Development* 135: 1813-1822.

CHROMOSOMAL LOCATION

Genetic locus: Galnt18 (mouse) mapping to 7 F1.

PRODUCT

GalNAc-TL4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GalNAc-TL4 shRNA Plasmid (m): sc-145315-SH and GalNAc-TL4 shRNA (m) Lentiviral Particles: sc-145315-V as alternate gene silencing products.

For independent verification of GalNAc-TL4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145315A, sc-145315B and sc-145315C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GalNAc-TL4 siRNA (m) is recommended for the inhibition of GalNAc-TL4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GalNAc-TL4 gene expression knockdown using RT-PCR Primer: GalNAc-TL4 (m)-PR: sc-145315-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.