

# GPR135 siRNA (m): sc-145702

## BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR135 (G protein-coupled receptor 135) is a 494 amino acid multi-pass membrane protein that belongs to the G protein-coupled receptor 1 family. Expressed in brain, eye, testis, cervix and testis, GPR135 shares high sequence homology with mouse and rat GPR135. The gene encoding GPR135 maps to human chromosome 14q23.1 and mouse chromosome 12 C3.

## REFERENCES

1. Takeda, S., Kadowaki, S., Haga, T., Takaesu, H. and Mitaku, S. 2002. Identification of G protein-coupled receptor genes from the human genome sequence. *FEBS Lett.* 520: 97-101.
2. Fredriksson, R., Höglund, P.J., Gloriam, D.E., Lagerstrom, M.C. and Schiöth, H.B. 2003. Seven evolutionarily conserved human rhodopsin G protein-coupled receptors lacking close relatives. *FEBS Lett.* 554: 381-388.
3. Vassilatis, D.K., Hohmann, J.G., Zeng, H., Li, F., Ranchalis, J.E., Mortrud, M.T., Brown, A., Rodriguez, S.S., Weller, J.R., Wright, A.C., Bergmann, J.E. and Gaitanaris, G.A. 2003. The G protein-coupled receptor repertoires of human and mouse. *Proc. Natl. Acad. Sci. USA* 100: 4903-4908.
4. Kristiansen, K. 2004. Molecular mechanisms of ligand binding, signaling, and regulation within the superfamily of G protein-coupled receptors: molecular modeling and mutagenesis approaches to receptor structure and function. *Pharmacol. Ther.* 103: 21-80.
5. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 607970. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Luttrell, L.M. 2006. Transmembrane signaling by G protein-coupled receptors. *Methods Mol. Biol.* 332: 3-49.

## CHROMOSOMAL LOCATION

Genetic locus: Gpr135 (mouse) mapping to 12 C3.

## PRODUCT

GPR135 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR135 shRNA Plasmid (m): sc-145702-SH and GPR135 shRNA (m) Lentiviral Particles: sc-145702-V as alternate gene silencing products.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

GPR135 siRNA (m) is recommended for the inhibition of GPR135 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR135 gene expression knockdown using RT-PCR Primer: GPR135 (m)-PR: sc-145702-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.