

Gstm5 siRNA (m): sc-145813

BACKGROUND

Members of the glutathione S-transferase (GST) family of proteins function in the detoxification of xenobiotics to protect cells against toxicant-induced damage. There are eight families of GST proteins, namely α , ζ , θ , κ , μ , π , σ and ω , each of which are composed of proteins that have a variety of functions throughout the cell. The GSTM proteins (GSTM1-GSTM5 in human and GSTM1-GSTM7 in mouse) are members of the μ class of enzymes that conjugate with glutathione and function in the detoxification of carcinogens, environmental toxins and products of oxidative stress. GSTM5 (glutathione S-transferase μ 5), also designated GTM5, is a 218 amino acid cytoplasmic protein belonging to the μ family and GST superfamily. Expressed in brain, lung and testis, GSTM5 is found at low levels in heart, and contains single N- and C-terminal GST domains.

REFERENCES

- Islam, M.Q., Platz, A., Szpirer, J., Szpirer, C., Levan, G. and Mannervik, B. 1989. Chromosomal localization of human glutathione transferase genes of classes α , μ and π . *Hum. Genet.* 82: 338-342.
- Pearson, W.R., Vorachek, W.R., Xu, S.J., Berger, R., Hart, I., Vannais, D. and Patterson, D. 1993. Identification of class- μ glutathione transferase genes GSTM1-GSTM5 on human chromosome 1p13. *Am. J. Hum. Genet.* 53: 220-233.
- Takahashi, Y., Campbell, E.A., Hirata, Y., Takayama, T. and Listowsky, I. 1993. A basis for differentiating among the multiple human μ -glutathione S-transferases and molecular cloning of brain GSTM5. *J. Biol. Chem.* 268: 8893-8898.
- Xu, S., Wang, Y., Roe, B. and Pearson, W.R. 1998. Characterization of the human class μ glutathione S-transferase gene cluster and the GSTM1 deletion. *J. Biol. Chem.* 273: 3517-3527.
- Delles, C., Padmanabhan, S., Lee, W.K., Miller, W.H., McBride, M.W., McClure, J.D., Brain, N.J., Wallace, C., Marcano, A.C., Schmieder, R.E., Brown, M.J., Caulfield, M.J., Munroe, P.B., Farrall, M., et al. 2008. Glutathione S-transferase variants and hypertension. *J. Hypertens.* 26: 1343-1352.
- Breton, C.V., Vora, H., Salam, M.T., Islam, T., Wenten, M., Gauderman, W.J., Van den Berg, D., Berhane, K., Peters, J.M. and Gilliland, F.D. 2009. Variation in the GST μ locus and tobacco smoke exposure as determinants of childhood lung function. *Am. J. Respir. Crit. Care Med.* 179: 601-607.
- Yu, K.D., Fan, L., Di, G.H., Yuan, W.T., Zheng, Y., Huang, W., Chen, A.X., Yang, C., Wu, J., Shen, Z.Z. and Shao, Z.M. 2009. Genetic variants in GSTM3 gene within GSTM4-GSTM2-GSTM1-GSTM5-GSTM3 cluster influence breast cancer susceptibility depending on GSTM1. *Breast Cancer Res. Treat.* 121: 485-496.
- Online Mendelian Inheritance in Man, OMIM[™]. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 138385. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Gstm5 (mouse) mapping to 3 F2.3.

PRODUCT

Gstm5 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Gstm5 shRNA Plasmid (m): sc-145813-SH and Gstm5 shRNA (m) Lentiviral Particles: sc-145813-V as alternate gene silencing products.

For independent verification of Gstm5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145813A and sc-145813B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Gstm5 siRNA (m) is recommended for the inhibition of Gstm5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Gstm5 gene expression knockdown using RT-PCR Primer: Gstm5 (m)-PR: sc-145813-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.