

Gucy2g siRNA (m): sc-145839

BACKGROUND

Guanylate cyclases belong to the adenylyl cyclase class-4/guanylyl cyclase family. There are two forms of guanylate cyclase. The soluble form, known as GCS or sGC, act as a receptor for nitric oxide. The membrane-bound receptor form, known as GC, is a peptide hormone receptor. GCS is a cGMP-synthesizing enzyme, which is the major receptor for the neurotransmitter nitric oxide (NO). It plays a crucial role in smooth muscle contractility, platelet reactivity and neurotransmission. The intracellular stimulation of guanylate cyclase by calcium, a key event in the recovery of the dark state of rod photoreceptors after exposure to light, is mediated by guanylate cyclase-activating protein (GCAP1). GCAPs are calcium-binding proteins belonging to the calmodulin superfamily. Gucy2g (guanylate cyclase 2g) is a 1,100 amino acid protein that is encoded by a gene that maps to mouse chromosome 19 D2.

REFERENCES

1. Yuen, P., Potter, L. and Garbers, D. 1990. A new form of guanylyl cyclase is preferentially expressed in rat kidney. *Biochemistry* 29: 10872-10878.
2. Wedel, B., Harteneck, C., Foerster, J., Friebe, A., Schultz, G. and Koesling, D. 1995. Functional domains of soluble guanylyl cyclase. *J. Biol. Chem.* 270: 24871-24875.
3. Bellamy, T., Wood, J., Goodwin, D. and Farthwaite, J. 2000. Rapid desensitization of the nitric oxide receptor, soluble guanylyl cyclase, underlies diversity of cellular cGMP responses. *Proc. Natl. Acad. Sci. USA* 97: 2928-2933.
4. Lee, Y., Martin, E. and Murad, F. 2000. Human recombinant soluble guanylyl cyclase: expression, purification and regulation. *Proc. Nat. Acad. Sci. USA* 97: 10763-10768.
5. Ibarra, C., Nedvetsky, P., Gerlach, M., Riederer, P. and Schmidt, H. 2001. Regional and age-dependent expression of the nitric oxide receptor, soluble guanylyl cyclase, in the human brain. *Brain Res.* 907: 54-60.
6. Koblin, M., Vehse, K., Budaus, L., Scholz, H. and Behrends, S. 2001. Nitric oxide activates the β 2 subunit of soluble guanylyl cyclase in the absence of a second subunit. *J. Biol. Chem.* 276: 30737-30743.
7. Martin, E., Le, Y. and Murad, F. 2001. YC-1 activation of human soluble guanylyl cyclase has both heme-dependent and heme independent components. *Proc. Natl. Acad. Sci. USA* 98: 12938-12942.
8. Gibb, B. and Garthwaite, J. 2001. Subunits of nitric oxide receptor, soluble guanylyl cyclase, expressed in rat brain. *Eur. J. Neurosci.* 13: 539-544.

CHROMOSOMAL LOCATION

Genetic locus: Gucy2g (mouse) mapping to 19 D2.

PRODUCT

Gucy2g siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Gucy2g shRNA Plasmid (m): sc-145839-SH and Gucy2g shRNA (m) Lentiviral Particles: sc-145839-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Gucy2g siRNA (m) is recommended for the inhibition of Gucy2g expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Gucy2g gene expression knockdown using RT-PCR Primer: Gucy2g (m)-PR: sc-145839-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.