

HEATR2 siRNA (m): sc-145921

BACKGROUND

HEATR2 (HEAT repeat containing 2) is an 855 amino acid protein that exists as three alternatively spliced isoforms and contains ten HEAT (Huntingtin, elongation factor 3 (EF3), protein phosphatase 2A (PP2A) and the yeast PI3-kinase Tor1) repeats. HEAT repeats form rod-like helical structures that are involved in intracellular transport. HEATR2 is encoded by a gene located on human chromosome 7p22.3. Human chromosome 7 houses over 1,000 genes and comprises nearly 5% of the human genome. Defects in some of the genes localized to chromosome 7 have been linked to Osteogenesis imperfecta, Williams-Beuren syndrome, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome.

REFERENCES

1. Andrade, M.A. and Bork, P. 1995. HEAT repeats in the Huntington's disease protein. *Nat. Genet.* 11: 115-116.
2. Seroussi, E., Pan, H.Q., Kedra, D., Roe, B.A. and Dumanski, J.P. 1998. Characterization of the human NIPSNAP1 gene from 22q12: a member of a novel gene family. *Gene* 212: 13-20.
3. Wang, X.Y., Smith, D.I., Liu, W. and James, C.D. 1998. GBAS, a novel gene encoding a protein with tyrosine phosphorylation sites and a transmembrane domain, is co-amplified with EGFR. *Genomics* 49: 448-451.
4. Groves, M.R., Hanlon, N., Turowski, P., Hemmings, B.A. and Barford, D. 1999. The structure of the protein phosphatase 2A PR65/A subunit reveals the conformation of its 15 tandemly repeated HEAT motifs. *Cell* 96: 99-110.
5. Lopez-Gines, C., Cerda-Nicolas, M., Gil-Benso, R., Pellin, A., Lopez-Guerrero, J.A., Callaghan, R., Benito, R., Roldan, P., Piquer, J., Llacer, J. and Barbera, J. 2005. Association of chromosome 7, chromosome 10 and EGFR gene amplification in glioblastoma multiforme. *Clin. Neuropathol.* 24: 209-218.
6. Ruano, Y., Mollejo, M., Ribalta, T., Fiaño, C., Camacho, F.I., Gómez, E., de Lope, A.R., Hernández-Moneo, J.L., Martínez, P. and Melendez, B. 2006. Identification of novel candidate target genes in amplicons of Glioblastoma multiforme tumors detected by expression and CGH microarray profiling. *Mol. Cancer* 5: 39.
7. Lo, K.C., Rossi, M.R., LaDuca, J., Hicks, D.G., Turpaz, Y., Hawthorn, L. and Cowell, J.K. 2007. Candidate glioblastoma development gene identification using concordance between copy number abnormalities and gene expression level changes. *Genes Chromosomes Cancer* 46: 875-894.
8. Necesalová, E., Vranová, V., Kuglík, P., Cejpek, P., Jarosová, M., Pesáková, M., Relichová, J. and Veselská, R. 2007. Incidence of the main genetic markers in glioblastoma multiforme is independent of tumor topology. *Neoplasma* 54: 212-218.
9. Smits, P., Rodenburg, R.J., Smeitink, J.A. and van den Heuvel, L.P. 2009. Sequence variants in four candidate genes (NIPSNAP1, GBAS, CHCHD1 and METT11D1) in patients with combined oxidative phosphorylation system deficiencies. *J. Inherit. Metab. Dis.* 3: S13-S19.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: *Heatr2* (mouse) mapping to 5 G2.

PRODUCT

HEATR2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HEATR2 shRNA Plasmid (m): sc-145921-SH and HEATR2 shRNA (m) Lentiviral Particles: sc-145921-V as alternate gene silencing products.

For independent verification of HEATR2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145921A, sc-145921B and sc-145921C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HEATR2 siRNA (m) is recommended for the inhibition of HEATR2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HEATR2 gene expression knockdown using RT-PCR Primer: HEATR2 (m)-PR: sc-145921-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.