



HEM1 siRNA (m): sc-145935

BACKGROUND

HEM1 (hematopoietic protein 1), also known as NCKAP1L (NCK-associated protein 1-like), is a 1,127 amino acid single-pass membrane protein that localizes to the cytoplasmic side of the cell membrane. One of several members of the highly conserved HEM family of tissue-specific transmembrane proteins, HEM1 is expressed in cells of hematopoietic origin where it is thought to play an important role in oogenesis. The gene encoding HEM1 maps to human chromosome 12q13.13, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and Trisomy 12p, which causes facial developmental defects and seizure disorders.

REFERENCES

1. Hromas, R., Collins, S., Raskind, W., Deaven, L. and Kaushansky, K. 1991. HEM1, a potential membrane protein, with expression restricted to blood cells. *Biochim. Biophys. Acta* 1090: 241-244.
2. Online Mendelian Inheritance in Man, OMIM™. 1993. Johns Hopkins University, Baltimore, MD. MIM Number: 141180. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Baumgartner, S., Martin, D., Chiquet-Ehrismann, R., Sutton, J., Desai, A., Huang, I., Kato, K. and Hromas, R. 1995. The HEM proteins: a novel family of tissue-specific transmembrane proteins expressed from invertebrates through mammals with an essential function in oogenesis. *J. Mol. Biol.* 251: 41-49.
4. Weiner, O.D., Rentel, M.C., Ott, A., Brown, G.E., Jedrychowski, M., Yaffe, M.B., Gygi, S.P., Cantley, L.C., Bourne, H.R. and Kirschner, M.W. 2006. HEM1 complexes are essential for Rac activation, Actin polymerization, and myosin regulation during neutrophil chemotaxis. *PLoS Biol.* 4: e38.
5. Joshi, A.D., Hegde, G.V., Dickinson, J.D., Mittal, A.K., Lynch, J.C., Eudy, J.D., Armitage, J.O., Bierman, P.J., Bociek, R.G., Devetten, M.P., Vose, J.M. and Joshi, S.S. 2007. Atm, CTLA4, MND4, and HEM1 in high versus low CD38 expressing B-cell chronic lymphocytic leukemia. *Clin. Cancer Res.* 13: 5295-5304.
6. Weiner, O.D., Marganski, W.A., Wu, L.F., Altschuler, S.J. and Kirschner, M.W. 2007. An Actin-based wave generator organizes cell motility. *PLoS Biol.* 5: e221.

CHROMOSOMAL LOCATION

Genetic locus: Nckap1l (mouse) mapping to 15 F3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

HEM1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HEM1 shRNA Plasmid (m): sc-145935-SH and HEM1 shRNA (m) Lentiviral Particles: sc-145935-V as alternate gene silencing products.

For independent verification of HEM1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145935A, sc-145935B and sc-145935C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HEM1 siRNA (m) is recommended for the inhibition of HEM1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HEM1 gene expression knockdown using RT-PCR Primer: HEM1 (m)-PR: sc-145935-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.