

HS3ST6 siRNA (m): sc-146088

BACKGROUND

Heparan sulfate structures, which are responsible for executing multiple biological activities, are generated and regulated by heparan sulfate biosynthetic enzymes. HS3ST6 (heparan sulfate glucosamine 3-O-sulfotransferase 6) is a 342 amino acid single-pass type II transmembrane protein that localizes to the golgi apparatus and belongs to the sulfotransferase 1 family. HS3ST6 transfers sulfate to the 3-OH position of the glucosamine residue of heparan sulfate to form 3-O-sulfated heparan sulfate. Due to observed susceptibility of HS3ST6-transfected CHO cells to HSV-1 infection, it has been suggested that HS3ST6 produces a specific entry receptor for HSV-1. The gene encoding HS3ST6 maps to human chromosome 16p13.3, which encodes over 900 genes and comprises nearly 3% of the human genome.

REFERENCES

1. Nakato, H. and Kimata, K. 2002. Heparan sulfate fine structure and specificity of proteoglycan functions. *Biochim. Biophys. Acta* 1573: 312-318.
2. Kamimura, K., Rhodes, J.M., Ueda, R., McNeely, M., Shukla, D., Kimata, K., Spear, P.G., Shworak, N.W. and Nakato, H. 2004. Regulation of Notch signaling by *Drosophila* heparan sulfate 3-O sulfotransferase. *J. Cell Biol.* 166: 1069-1079.
3. Xu, D., Tiwari, V., Xia, G., Clement, C., Shukla, D. and Liu, J. 2005. Characterization of heparan sulphate 3-O-sulphotransferase isoform 6 and its role in assisting the entry of herpes simplex virus type 1. *Biochem. J.* 385: 451-459.
4. Muñoz, E., Xu, D., Kemp, M., Zhang, F., Liu, J. and Linhardt, R.J. 2006. Affinity, kinetic, and structural study of the interaction of 3-O-sulfotransferase isoform 1 with heparan sulfate. *Biochemistry* 45: 5122-5128.
5. de Agostini, A.I., Dong, J.C., de Vantery Arrighi, C., Ramus, M.A., Dentand-Quadri, I., Thalmann, S., Ventura, P., Ibecheole, V., Monge, F., Fischer, A.M., HajMohammadi, S., Shworak, N.W., et al. 2008. Human follicular fluid heparan sulfate contains abundant 3-O-sulfated chains with anticoagulant activity. *J. Biol. Chem.* 283: 28115-28124.
6. Mantelli, F., Schaffer, L., Dana, R., Head, S.R. and Argüeso, P. 2009. Glycogene expression in conjunctiva of patients with dry eye: downregulation of Notch signaling. *Invest. Ophthalmol. Vis. Sci.* 50: 2666-2672.

CHROMOSOMAL LOCATION

Genetic locus: Hs3st6 (mouse) mapping to 17 A3.3.

PRODUCT

HS3ST6 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HS3ST6 shRNA Plasmid (m): sc-146088-SH and HS3ST6 shRNA (m) Lentiviral Particles: sc-146088-V as alternate gene silencing products.

For independent verification of HS3ST6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146088A and sc-146088B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HS3ST6 siRNA (m) is recommended for the inhibition of HS3ST6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HS3ST6 gene expression knockdown using RT-PCR Primer: HS3ST6 (m)-PR: sc-146088-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.