

IFI-44L siRNA (m): sc-146152

BACKGROUND

Interferon-induced protein 44-like (IFI-44L), also known as C1orf29, is a 452 amino acid cytoplasmic protein that shares some sequence similarities with IFI-44. IFI-44 is a cytoplasmic protein that aggregates to form microtubule structures. The genes that encode IFI-44L and IFI-44 are located on chromosome 1, which is the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes Lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

1. Watson, M.L., Kingsmore, S.F., Johnston, G.I., Siegelman, M.H., Le Beau, M.M., Lemons, R.S., Bora, N.S., Howard, T.A., Weissman, I.L., McEver, R.P. and Seldin, M.F. 1990. Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263-272.
2. Takahashi, K., Kitamura, N., Shibui, T., Kamizono, M., Matsui, R., Yoshiyama, Y., Maeda, T., Kondo, J., Honda, Y. and Yamada, E. 1990. Cloning, sequencing and expression in *Escherichia coli* of cDNA for a non-A, non-B hepatitis-associated microtubular aggregates protein. *J. Gen. Virol.* 71: 2005-2011.
3. Kitamura, A., Takahashi, K., Okajima, A. and Kitamura, N. 1994. Induction of the human gene for p44, a hepatitis-C-associated microtubular aggregate protein, by interferon- α/β . *Eur. J. Biochem.* 224: 877-883.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610468. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Izmailova, E., Bertley, F.M., Huang, Q., Makori, N., Miller, C.J., Young, R.A. and Aldovini, A. 2003. HIV-1 Tat reprograms immature dendritic cells to express chemoattractants for activated T cells and macrophages. *Nat. Med.* 9: 191-197.
6. Weise, A., Starke, H., Mrasek, K., Claussen, U. and Liehr, T. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
7. Marzin, Y., Jamet, D., Douet-Guilbert, N., Morel, F., Le Bris, M.J., Morice, P., Abgrall, J.F., Berthou, C. and De Braekeleer, M. 2006. Chromosome 1 abnormalities in multiple myeloma. *Anticancer Res.* 26: 953-959.
8. Hallen, L.C., Burki, Y., Ebeling, M., Broger, C., Siegrist, F., Oroszlan-Szovik, K., Bohrmann, B., Certa, U. and Foser, S. 2007. Antiproliferative activity of the human IFN- α -inducible protein IFI44. *J. Interferon Cytokine Res.* 27: 675-680.

CHROMOSOMAL LOCATION

Genetic locus: H28 (mouse) mapping to 3 H3.

PRODUCT

IFI-44L siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IFI-44L shRNA Plasmid (m): sc-146152-SH and IFI-44L shRNA (m) Lentiviral Particles: sc-146152-V as alternate gene silencing products.

For independent verification of IFI-44L (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146152A, sc-146152B and sc-146152C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IFI-44L siRNA (m) is recommended for the inhibition of IFI-44L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IFI-44L gene expression knockdown using RT-PCR Primer: IFI-44L (m)-PR: sc-146152-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.