

IL-1R8 siRNA (m): sc-146213

BACKGROUND

IL-1R8, also known as X-linked interleukin-1 receptor accessory protein-like 1 (IL1RAPL-1), oligophrenin-4 or three immunoglobulin domain-containing IL-1 receptor-related 2 (TIGIRR-2), is a member of the interleukin-1/Toll-like receptor family predominantly expressed in brain and heart. It is a single-pass transmembrane protein with a highly conserved cytoplasmic region that is slightly longer than that of other family members. IL-1R8 is highly homologous to IL-1R9 and the IL-1 accessory protein receptor chains. The genes encoding IL-1R8 and IL-1R9 are both found on the X chromosome. IL-1R8 may interact with NCS-1 and, via this interaction, may be involved in the regulation of exocytosis and synaptic transmission. A truncating mutation in IL-1R8, possibly disrupting the activity of N-type voltage-gated calcium channels or NCS-1 dependent neuronal activity, results in non-syndromic mental retardation X-linked type 21 or MRX21.

REFERENCES

1. Born, T.L., Smith, D.E., Garka, K.E., Renshaw, B.R., Bertles, J.S. and Sims, J.E. 2000. Identification and characterization of two members of a novel class of the interleukin-1 receptor (IL-1R) family. Delineation of a new class of IL-1R-related proteins based on signaling. *J. Biol. Chem.* 275: 29946-29954.
2. Bahi, N., Friocourt, G., Carrie, A., Graham, M.E., Weiss, J.L., Chafey, P., Fauchereau, F., Burgoyne, R.D. and Chelly, J. 2003. IL1 receptor accessory protein like, a protein involved in X-linked mental retardation, interacts with Neuronal Calcium Sensor-1 and regulates exocytosis. *Hum. Mol. Genet.* 12: 1415-1425.
3. Zhang, Y.H., Huang, B.L., Niakan, K.K., McCabe, L.L., McCabe, E.R. and Dipple, K.M. 2004. IL1RAPL1 is associated with mental retardation in patients with complex glycerol kinase deficiency who have deletions extending telomeric of DAX1. *Hum. Mutat.* 24: 273-273.
4. Dinarello, C.A. 2004. The IL-1 family and inflammatory diseases. *Clin. Exp. Rheumatol.* 20: S1-S13.
5. Andre, R., Lerouet, D., Kimber, I., Pinteaux, E. and Rothwell, N.J. 2005. Regulation of expression of the novel IL-1 receptor family members in the mouse brain. *J. Neurochem.* 95: 324-330.
6. Tabolacci, E., Pomponi, M.G., Pietrobono, R., Terracciano, A., Chiurazzi, P. and Neri, G. 2006. A truncating mutation in the IL1RAPL1 gene is responsible for X-linked mental retardation in the MRX21 family. *Am. J. Med. Genet. A* 140: 482-487.
7. Gambino, F., Pavlowsky, A., Begle, A., Dupont, J.L., Bahi, N., Courjaret, R., Gardette, R., Hadjkacem, H., Skala, H., Poulain, B., Chelly, J., Vitale, N. and Humeau, Y. 2007. IL1-receptor accessory protein-like 1 (IL1RAPL1), a protein involved in cognitive functions, regulates N-type Ca²⁺-channel and neurite elongation. *Proc. Natl. Acad. Sci. USA* 104: 9063-9068.
8. Barbaro, M., Oscarson, M., Schoumans, J., Staaf, J., Ivarsson, S.A. and Wedell, A. 2007. Isolated 46,XY gonadal dysgenesis in two sisters caused by a Xp21.2 interstitial duplication containing the DAX1 gene. *J. Clin. Endocrinol. Metab.* 92: 3305-3313.

CHROMOSOMAL LOCATION

Genetic locus: IL1rapl1 (mouse) mapping to X C1.

PRODUCT

IL-1R8 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-1R8 shRNA Plasmid (m): sc-146213-SH and IL-1R8 shRNA (m) Lentiviral Particles: sc-146213-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IL-1R8 siRNA (m) is recommended for the inhibition of IL-1R8 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-1R8 gene expression knockdown using RT-PCR Primer: IL-1R8 (m)-PR: sc-146213-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.