



KIF9 siRNA (m): sc-146478

BACKGROUND

Kinesin is a cytoskeletal motor protein involved in axonal transport and cell division. The kinesins constitute a large family of microtubule-dependent motor proteins, which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual kinesin members play crucial roles in cell division, intracellular transport and membrane trafficking events, including endocytosis and transcytosis. KIF9 (kinesin family member 9) is a member of the kinesin-like protein family and is a 790 amino acid protein that exists as two isoforms. KIF9 contains one kinesin-motor domain and KIF9's expression is developmentally regulated in tissues including brain, kidney, spleen, lung and testis. KIF9 is involved in keeping the MTOC (microtubule organizing center) connected to the nucleus during interphase, and is thought to interact with Gem, an association which may connect KIF9 to the cytoskeleton. In addition, KIF9 may act as a plus-ended microtubule motor that may exist as a homodimer.

REFERENCES

1. Nakagawa, T., Tanaka, Y., Matsuoka, E., Kondo, S., Okada, Y., Noda, Y., Kanai, Y. and Hirokawa, N. 1997. Identification and classification of 16 new kinesin superfamily (KIF) proteins in mouse genome. *Proc. Natl. Acad. Sci. USA* 94: 9654-9659.
2. Piddini, E., Schmid, J.A., de Martin, R. and Dotti, C.G. 2001. The Ras-like GTPase Gem is involved in cell shape remodelling and interacts with the novel kinesin-like protein KIF9. *EMBO J.* 20: 4076-4087.
3. Miki, H., Setou, M., Kaneshiro, K. and Hirokawa, N. 2001. All kinesin superfamily protein, KIF, genes in mouse and human. *Proc. Natl. Acad. Sci. USA* 98: 7004-7011.
4. Miki, H., Setou, M. and Hirokawa, N. 2003. Kinesin superfamily proteins (KIFs) in the mouse transcriptome. *Genome Res.* 13: 1455-1465.
5. Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607910. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Takano, K., Miki, T., Katahira, J. and Yoneda, Y. 2007. NXF2 is involved in cytoplasmic mRNA dynamics through interactions with motor proteins. *Nucleic Acids Res.* 35: 2513-2521.
7. Tikhonenko, I., Nag, D.K., Robinson, D.N. and Koonce, M.P. 2009. Microtubule-nucleus interactions in *Dictyostelium discoideum* mediated by central motor kinesins. *Eukaryotic Cell* 8: 723-731.

CHROMOSOMAL LOCATION

Genetic locus: Kif9 (mouse) mapping to 9 F2.

PRODUCT

KIF9 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KIF9 shRNA Plasmid (m): sc-146478-SH and KIF9 shRNA (m) Lentiviral Particles: sc-146478-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KIF9 siRNA (m) is recommended for the inhibition of KIF9 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KIF9 gene expression knockdown using RT-PCR Primer: KIF9 (m)-PR: sc-146478-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.