

KLHL4 siRNA (m): sc-146533

BACKGROUND

The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (Poxvirus and zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C₂H₂-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. KLHL4 (kelch-like 4), also known as DKELCHL, is a 718 amino acid cytoplasmic protein that is expressed in adult fibroblasts and in a range of fetal tissues including tongue, palate and mandible. KLHL4 contains one BTB (POZ) domain and six Kelch repeats. KLHL4 exists as four alternatively spliced isoforms and is encoded by a gene located on human chromosome Xq21.31.

REFERENCES

1. Albagli, O., Dhordain, P., Deweindt, C., Lecocq, G. and Leprince, D. 1995. The BTB/POZ domain: a new protein-protein interaction motif common to DNA- and Actin-binding proteins. *Cell Growth Differ.* 6: 1193-1198.
2. Adams, J., Kelso, R. and Cooley, L. 2000. The kelch repeat superfamily of proteins: propellers of cell function. *Trends Cell Biol.* 10: 17-24.
3. Braybrook, C., Warry, G., Howell, G., Arnason, A., Bjornsson, A., Moore, G.E., Ross, M.T. and Stanier, P. 2001. Identification and characterization of KLHL4, a novel human homologue of the *Drosophila* Kelch gene that maps within the X-linked cleft palate and Ankyloglossia (CPX) critical region. *Genomics* 72: 128-136.
4. Braybrook, C., Warry, G., Howell, G., Mandryko, V., Arnason, A., Bjornsson, A., Ross, M.T., Moore, G.E. and Stanier, P. 2001. Physical and transcriptional mapping of the X-linked cleft palate and ankyloglossia (CPX) critical region. *Hum. Genet.* 108: 537-545.
5. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 300348. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Prag, S. and Adams, J.C. 2003. Molecular phylogeny of the kelch-repeat superfamily reveals an expansion of BTB/Kelch proteins in animals. *BMC Bioinformatics* 4: 42.
7. Stogios, P.J. and Prive, G.G. 2004. The BACK domain in BTB-kelch proteins. *Trends Biochem. Sci.* 29: 634-637.

CHROMOSOMAL LOCATION

Genetic locus: Klhl4 (mouse) mapping to X E1.

PRODUCT

KLHL4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KLHL4 shRNA Plasmid (m): sc-146533-SH and KLHL4 shRNA (m) Lentiviral Particles: sc-146533-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KLHL4 siRNA (m) is recommended for the inhibition of KLHL4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KLHL4 gene expression knockdown using RT-PCR Primer: KLHL4 (m)-PR: sc-146533-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.