

# Kremen-2 siRNA (m): sc-146568

## BACKGROUND

Kremen-2, also known as KRM2, is a 462 amino acid single-pass type I membrane protein that interacts with XTP3-B. Existing as four alternatively spliced isoforms, Kremen-2 contains one CUB domain, one kringle domain and one WSC domain. As a receptor for Dickkopf (Dkk-1) protein, Kremen-2 cooperates with Dkk-1 to block Wnt/ $\beta$ -catenin signaling. Kremen-2 forms a ternary complex with Dkk1 and LRP6 to induce rapid endocytosis and removal of the Wnt receptor LRP6 from the plasma membrane. The gene that encodes Kremen-2 consists of approximately 4,222 bases and maps to human chromosome 16p13.3. Encoding over 900 genes and consisting of approximately 90 million base pairs, chromosome 16 makes up nearly 3% of the human genome and is associated with a variety of genetic disorders, such as giant axonal neuropathy, Rubinstein-Taybi syndrome and Croh's disease.

## REFERENCES

1. Mentzer, W.C., Johnston, R.B., Baehner, R.L. and Nathan, D.G. 1977. An unusual form of chronic neutropenia in a father and daughter with hypogammaglobulinaemia. *Br. J. Haematol.* 36: 313-322.
2. Baraitser, M. and Preece, M.A. 1983. The Rubinstein-Taybi syndrome: occurrence in two sets of identical twins. *Clin. Genet.* 23: 318-320.
3. Breuning, M.H., Dauwerse, H.G., Fugazza, G., Saris, J.J., Spruit, L., Wijnen, H., Tommerup, N., van der Hagen, C.B., Imaizumi, K., Kuroki, Y., van den Boogaard, M.J., de Pater, J.M., Mariman, E.C., et al. 1993. Rubinstein-Taybi syndrome caused by submicroscopic deletions within 16p13.3. *Am. J. Hum. Genet.* 52: 249-254.
4. Mao, B., Wu, W., Davidson, G., Marhold, J., Li, M., Mechler, B.M., Delius, H., Hoppe, D., Stanek, P., Walter, C., Glinka, A. and Niehrs, C. 2002. Kremen proteins are Dickkopf receptors that regulate Wnt/ $\beta$ -catenin signalling. *Nature* 417: 664-667.
5. Kuhlensäumer, G., Young, P., Oberwittler, C., Hünermund, G., Schirmacher, A., Domschke, K., Ringelstein, B. and Stögbauer, F. 2002. Giant axonal neuropathy (GAN): case report and two novel mutations in the gigaxonin gene. *Neurology* 58: 1273-1276.
6. Cho, J.H. 2004. Advances in the genetics of inflammatory bowel disease. *Curr. Gastroenterol. Rep.* 6: 467-473.
7. Mathew, C.G. and Lewis, C.M. 2004. Genetics of inflammatory bowel disease: progress and prospects. *Hum. Mol. Genet.* 1: R161-R168.
8. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 609899. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
9. Dun, X., Jiang, H., Zou, J., Shi, J., Zhou, L., Zhu, R. and Hou, J. 2010. Differential expression of DKK-1 binding receptors on stromal cells and myeloma cells results in their distinct response to secreted DKK-1 in myeloma. *Mol. Cancer* 9: 247.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Kremen2 (mouse) mapping to 17 A3.3.

## PRODUCT

Kremen-2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Kremen-2 shRNA Plasmid (m): sc-146568-SH and Kremen-2 shRNA (m) Lentiviral Particles: sc-146568-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Kremen-2 siRNA (m) is recommended for the inhibition of Kremen-2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Kremen-2 gene expression knockdown using RT-PCR Primer: Kremen-2 (m)-PR: sc-146568-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.