

# KY peptidase siRNA (m): sc-146614

## BACKGROUND

Filamins are Actin-binding proteins which contain an N-terminal Actin-binding domain, a membrane glycoprotein domain and a C-terminal self-association domain. Filamins help reshape the cytoskeleton by forming flexible cross-links between two Actin filaments, which maintain membrane integrity during force application. Filamin 2, also designated Filamin C, is a skeletal- and cardiac-muscle specific form of Filamin, which binds  $\gamma$ -sarcoglycan and  $\delta$ -sarcoglycan, but not  $\alpha$ -sarcoglycan or  $\beta$ -sarcoglycan. KY peptidase (kyphoscoliosis peptidase) is a 561 amino acid cytoskeleton protease that interacts with several sarcomeric cytoskeletal proteins, including Filamin 2. KY peptidase probably plays a role in the maturation, function and stabilization of the neuromuscular junction. KY-null mouse mutants exhibit distinct irregular subcellular Filamin 2 localization, suggesting that KY peptidase deficiency may be the cause of several types of limb-girdle muscular dystrophies.

## REFERENCES

1. Blanco, G., Nikitopoulou, A., Kraus, M., Mason, R.M., Coulton, G.R. and Brown, S.D. 1998. A STS content physical and transcription map across the KY, kyphoscoliosis, nonrecombinant region. *Genomics* 54: 415-423.
2. van der Ven, P.F., Obermann, W.M., Lemke, B., Gautel, M., Weber, K. and Fürst, D.O. 2000. Characterization of muscle filamin isoforms suggests a possible role of  $\gamma$ -Filamin/ABP-L in sarcomeric Z-disc formation. *Cell Motil. Cytoskeleton* 45: 149-162.
3. Thompson, T.G., Chan, Y.M., Hack, A.A., Brosius, M., Rajala, M., Lidov, H.G., McNally, E.M., Watkins, S. and Kunkel, L.M. 2000. Filamin 2 (FLN2): a muscle-specific sarcoglycan interacting protein. *J. Cell Biol.* 148: 115-126.
4. Blanco, G., Coulton, G.R., Biggin, A., Grainge, C., Moss, J., Barrett, M., Berquin, A., Marechal, G., Skynner, M., van Mier, P., Nikitopoulou, A., Kraus, M., Ponting, C.P., Mason, R.M. and Brown, S.D. 2001. The kyphoscoliosis (KY) mouse is deficient in hypertrophic responses and is caused by a mutation in a novel muscle-specific protein. *Hum. Mol. Genet.* 10: 9-16.
5. Murray, J.T., Campbell, D.G., Pegg, M., Mora, A., Alfonso, M. and Cohen, P. 2004. Identification of Filamin C as a new physiological substrate of PKB $\alpha$  using KESTREL. *Biochem. J.* 384: 489-494.
6. Beatham, J., Romero, R., Townsend, S.K., Hacker, T., van der Ven, P.F. and Blanco, G. 2004. Filamin C interacts with the muscular dystrophy KY protein and is abnormally distributed in mouse KY deficient muscle fibres. *Hum. Mol. Genet.* 13: 2863-2874.
7. Anastasi, G., Cutroneo, G., Trimarchi, F., Santoro, G., Bruschetta, D., Bramanti, P., Pisani, A. and Favaloro, A. 2004. Evaluation of sarcoglycans, vinculin-talin-integrin system and Filamin 2 in  $\alpha$ - and  $\gamma$ -sarcoglycanopathy: an immunohistochemical study. *Int. J. Mol. Med.* 14: 989-999.
8. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 605739. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: Ky (mouse) mapping to 9 F1.

## PRODUCT

KY peptidase siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KY peptidase shRNA Plasmid (m): sc-146614-SH and KY peptidase shRNA (m) Lentiviral Particles: sc-146614-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

KY peptidase siRNA (m) is recommended for the inhibition of KY peptidase expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KY peptidase gene expression knockdown using RT-PCR Primer: KY peptidase (m)-PR: sc-146614-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.