

LEPROTL1 siRNA (m): sc-146710

BACKGROUND

Mutation of Ob (obesity factor), also known as leptin precursor, results in profound obesity and type II diabetes as part of a syndrome that resembles morbid obesity in humans. The Ob gene product may function as a component of a signaling pathway in adipose tissue that functions to regulate body fat depot size. The leptin receptor, designated Ob-R, has been shown to be a single membrane-spanning receptor that most resembles the gp130 signal transducing component of the IL-6, G-CSF and LIF receptor. LEPROTL1 (leptin receptor overlapping transcript-like 1) is a 131 amino acid multi-pass membrane protein that is highly homologous to the leptin receptor gene-related protein, Ob-R. LEPROTL1 is widely expressed and contains a JAK binding site. The LEPROTL1 gene is located on chromosome 8p12, which is made up of nearly 146 million bases and encodes about 800 genes.

REFERENCES

- Huang, Y., et al. 2001. Cloning and characterization of a novel human leptin receptor overlapping transcript-like 1 gene (LEPROTL1). *Biochim. Biophys. Acta* 1517: 327-331.
- Kim, J.H., et al. 2008. Minimizing a QTL region for intramuscular fat content by characterizing the porcine phosphodiesterase 4B (PDE4B) gene. *BMB Rep.* 41: 466-471.
- Kurokawa, T., et al. 2008. Genomic characterization and tissue distribution of leptin receptor and leptin receptor overlapping transcript genes in the pufferfish, *Takifugu rubripes*. *Gen. Comp. Endocrinol.* 158: 108-114.
- Hansen, G.H., et al. 2008. Leptin and the obesity receptor (OB-R) in the small intestine and colon: a colocalization study. *J. Histochem. Cytochem.* 56: 677-685.
- Jardé, T., et al. 2008. Leptin and leptin receptor involvement in cancer development: a study on human primary breast carcinoma. *Oncol. Rep.* 19: 905-911.
- De Luis, D.A., et al. 2009. Leptin and obesity. *Minerva Med.* 100: 229-236.
- Koros, C., et al. 2009. Fat diet affects leptin receptor levels in the rat cerebellum. *Nutrition* 25: 85-87.

CHROMOSOMAL LOCATION

Genetic locus: Leprotl1 (mouse) mapping to 8 A4.

PRODUCT

LEPROTL1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LEPROTL1 shRNA Plasmid (m): sc-146710-SH and LEPROTL1 shRNA (m) Lentiviral Particles: sc-146710-V as alternate gene silencing products.

For independent verification of LEPROTL1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146710A, sc-146710B and sc-146710C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LEPROTL1 siRNA (m) is recommended for the inhibition of LEPROTL1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LEPROTL1 gene expression knockdown using RT-PCR Primer: LEPROTL1 (m)-PR: sc-146710-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Wu, S., et al. 2013. Increased expression of fibroblast growth factor 21 (FGF21) during chronic undernutrition causes growth hormone insensitivity in chondrocytes by inducing leptin receptor overlapping transcript (LEPROTL1) and leptin receptor overlapping transcript-like 1 (LEPROTL1) expression. *J. Biol. Chem.* 288: 27375-27383.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.