



Lipocalin-8 siRNA (m): sc-146752

BACKGROUND

The Lipocalin family is composed of structurally conserved hydrophobic ligand-binding proteins and is represented in all major taxonomic groups from prokaryotes to primates. Members of the lipocalin family are characterized by several common molecular-recognition properties: the ability to bind a range of small hydrophobic molecules, binding to specific cell-surface receptors and the formation of complexes with soluble macromolecules. Lipocalin-8, also known as LCN8, is a 175 amino acid protein that is predominantly expressed in epididymis, where it is thought to regulate epididymis-specific gene expression. Belonging to the calycin superfamily, Lipocalin-8 may play a role in sperm maturation. The gene encoding Lipocalin-8 maps to human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome.

REFERENCES

1. Flower, D.R. 1995. Multiple molecular recognition properties of the lipocalin protein family. *J. Mol. Recognit.* 8: 185-195.
2. Flower, D.R. 1996. The lipocalin protein family: structure and function. *Biochem. J.* 318: 1-14.
3. Suzuki, K., Lareyre, J.J., Sánchez, D., Gutierrez, G., Araki, Y., Matusik, R.J. and Orgebin-Crist, M.C. 2004. Molecular evolution of epididymal lipocalin genes localized on mouse chromosome 2. *Gene* 339: 49-59.
4. Humphray, S.J., Oliver, K., Hunt, A.R., Plumb, R.W., Loveland, J.E., Howe, K.L., Andrews, T.D., Searle, S., Hunt, S.E., Scott, C.E., Jones, M.C., Ainscough, R., Almeida, J.P., Ambrose, K.D., Ashwell, R.I., et al. 2004. DNA sequence and analysis of human chromosome 9. *Nature* 429: 369-374.
5. Grzyb, J., Latowski, D. and Strzałka, K. 2006. Lipocalins-a family portrait. *J. Plant Physiol.* 163: 895-915.
6. Suzuki, K., Yu, X., Chaurand, P., Araki, Y., Lareyre, J.J., Caprioli, R.M., Orgebin-Crist, M.C. and Matusik, R.J. 2007. Epididymis-specific lipocalin promoters. *Asian J. Androl.* 9: 515-521.

CHROMOSOMAL LOCATION

Genetic locus: Lcn8 (mouse) mapping to 2 A3.

PRODUCT

Lipocalin-8 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Lipocalin-8 shRNA Plasmid (m): sc-146752-SH and Lipocalin-8 shRNA (m) Lentiviral Particles: sc-146752-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Lipocalin-8 siRNA (m) is recommended for the inhibition of Lipocalin-8 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Lipocalin-8 gene expression knockdown using RT-PCR Primer: Lipocalin-8 (m)-PR: sc-146752-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.