

LSmD1 siRNA (m): sc-149134

BACKGROUND

Sm and Sm-like (LSm) proteins form donut shaped heptameric complexes that are involved in various steps of RNA metabolism. Lsm proteins facilitate RNA protein interactions and structural changes that are required during ribosomal subunit assembly. LSmD1 (LSM domain-containing protein 1), also known as MAK31 and PFAAP2 (phosphonoformate immuno-associated protein 2), is a 125 amino acid protein that is a component of the N-terminal acetyltransferase C (NatC) complex. Composed of MAK10, NAT-12 and LSmD1, the NatC complex catalyzes the acetylation of amino-terminal methionine residues. siRNA knockdown of NatC complex subunits leads to p53-dependent cell death and reduced growth of cell lines. There are two isoforms of LSmD1 that are produced as a result of alternative splicing events.

REFERENCES

1. Wickner, R.B., Fujimura, T. and Esteban, R. 1986. Overview of double-stranded RNA replication in *Saccharomyces cerevisiae*. Basic Life Sci. 40: 149-163.
2. Treich, I., Ho, L. and Carlson, M. 1998. Direct interaction between Rsc6 and Rsc8/Swh3, two proteins that are conserved in SWI/SNF-related complexes. Nucleic Acids Res. 26: 3739-3745.
3. Lanzuolo, C., Ederle, S., Pollice, A., Russo, F., Storlazzi, A. and Pulitzer, J.F. 2001. The HTL1 gene (YCR020W-b) of *Saccharomyces cerevisiae* is necessary for growth at 37 degrees C, and for the conservation of chromosome stability and fertility. Yeast 18: 1317-1330.
4. Polevoda, B. and Sherman, F. 2003. Composition and function of the eukaryotic N-terminal acetyltransferase subunits. Biochem. Biophys. Res. Commun. 308: 1-11.
5. Polevoda, B. and Sherman, F. 2003. N-terminal acetyltransferases and sequence requirements for N-terminal acetylation of eukaryotic proteins. J. Mol. Biol. 325: 595-622.
6. Wenzlau, J.M., Garl, P.J., Simpson, P., Stenmark, K.R., West, J., Artinger, K.B., Nemenoff, R.A. and Weiser-Evans, M.C. 2006. Embryonic growth-associated protein is one subunit of a novel N-terminal acetyltransferase complex essential for embryonic vascular development. Circ. Res. 98: 846-855.
7. Polevoda, B., Brown, S., Cardillo, T.S., Rigby, S. and Sherman, F. 2008. Yeast N^α-terminal acetyltransferases are associated with ribosomes. J. Cell. Biochem. 103: 492-508.
8. Starheim, K.K., Gromyko, D., Evjenth, R., Rynningen, A., Varhaug, J.E., Lillehaug, J.R. and Arnesen, T. 2009. Knockdown of human N^α-terminal acetyltransferase complex C leads to p53-dependent apoptosis and aberrant human Arl8b localization. Mol. Cell. Biol. 29: 3569-3581.

CHROMOSOMAL LOCATION

Genetic locus: LsmD1 (mouse) mapping to 11 B3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

LSmD1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LSmD1 shRNA Plasmid (m): sc-149134-SH and LSmD1 shRNA (m) Lentiviral Particles: sc-149134-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LSmD1 siRNA (m) is recommended for the inhibition of LSmD1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LSmD1 gene expression knockdown using RT-PCR Primer: LSmD1 (m)-PR: sc-149134-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.