

MIPU1 siRNA (m): sc-149441

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. MIPU1, also known as ZNF667 (zinc finger protein 667), is a 610 amino acid nuclear protein that is thought to play a role in transcriptional regulation and belongs to the Krüppel C₂H₂-type zinc-finger protein family. MIPU1 contains one KRAB domain and 15 C₂H₂-type zinc fingers, and is encoded by a gene that maps to human chromosome 19q13.43.

REFERENCES

1. Bray, P., Lichter, P., Thiesen, H.J., Ward, D.C. and Dawid, I.B. 1991. Characterization and mapping of human genes encoding zinc finger proteins. *Proc. Natl. Acad. Sci. USA* 88: 9563-9567.
2. Lichter, P., Bray, P., Ried, T., Dawid, I.B. and Ward, D.C. 1992. Clustering of C₂H₂ zinc finger motif sequences within telomeric and fragile site regions of human chromosomes. *Genomics* 13: 999-1007.
3. Matakai, C., Murakami, T., Umetani, M., Wada, Y., Ishii, M., Tsutsumi, S., Aburatani, H., Hamakubo, T. and Kodama, T. 2000. A novel zinc finger protein mRNA in human umbilical vein endothelial cells is profoundly induced by tumor necrosis factor α . *J. Atheroscler. Thromb.* 7: 97-103.
4. Urrutia, R. 2003. KRAB-containing zinc-finger repressor proteins. *Genome Biol.* 4: 231.
5. Huntley, S., Baggott, D.M., Hamilton, A.T., Tran-Gyamfi, M., Yang, S., Kim, J., Gordon, L., Branscomb, E. and Stubbs, L. 2006. A comprehensive catalog of human KRAB-associated zinc finger genes: insights into the evolutionary history of a large family of transcriptional repressors. *Genome Res.* 16: 669-677.
6. Filion, G.J., Zhenilo, S., Salozhin, S., Yamada, D., Prokhortchouk, E. and Defossez, P.A. 2006. A family of human zinc finger proteins that bind methylated DNA and repress transcription. *Mol. Cell. Biol.* 26: 169-181.
7. Tian, C.Y., Zhang, L.Q. and He, F.C. 2006. Progress in the study of KRAB zinc finger protein. *Yi Chuan* 28: 1451-1456.

CHROMOSOMAL LOCATION

Genetic locus: Zfp667 (mouse) mapping to 7 A1.

PRODUCT

MIPU1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MIPU1 shRNA Plasmid (m): sc-149441-SH and MIPU1 shRNA (m) Lentiviral Particles: sc-149441-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MIPU1 siRNA (m) is recommended for the inhibition of MIPU1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MIPU1 gene expression knockdown using RT-PCR Primer: MIPU1 (m)-PR: sc-149441-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.