

Mucin 3 siRNA (m): sc-149703

BACKGROUND

The Mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats (VNTRs) encoded by 60 base pairs (Mucin 1), 69 base pairs (Mucin 2) and 51 base pairs (Mucin 3). The number of repeats is highly polymorphic and varies among different alleles. Mucin 1 proteins are expressed as type I membrane proteins in addition to secreted forms. Mucin 1 is aberrantly expressed in epithelial tumors including breast carcinomas. Mucin 2 coats the epithelia of the intestines and airways and is associated with colonic tumors. Mucin 3 is a major component of various mucus gels and is broadly expressed in normal and tumor cells.

REFERENCES

1. Siddiqui, J., et al. 1988. Isolation and sequencing of a cDNA coding for the human DF3 breast carcinoma-associated antigen. *Proc. Natl. Acad. Sci. USA* 85: 2320-2323.
2. Lan, M.S., et al. 1990. Cloning and sequencing of a human pancreatic tumor Mucin cDNA. *J. Biol. Chem.* 265: 15294-15299.
3. Gum, J.R., et al. 1990. Molecular cloning of cDNAs derived from a novel human intestinal Mucin gene. *Biochem. Biophys. Res. Commun.* 171: 407-415.
4. Gum, J.R., et al. 1992. The human MUC2 intestinal Mucin has cysteine-rich subdomains located both upstream and downstream of its central repetitive region. *J. Biol. Chem.* 267: 21375-21383.
5. Pandey, P., et al. 1995. Association of the DF3/MUC1 breast cancer antigen with GRB2 and the Sos/Ras exchange protein. *Cancer Res.* 55: 4000-4003.
6. Geng, H., et al. 2006. Soluble form of T cell Ig Mucin 3 is an inhibitory molecule in T cell-mediated immune response. *J. Immunol.* 176: 1411-1420.
7. Frisnacho-Kiss, S., et al. 2006. Cutting edge: T cell Ig Mucin 3 reduces inflammatory heart disease by increasing CTLA-4 during innate immunity. *J. Immunol.* 176: 6411-6415.
8. Louis, N.A., et al. 2006. Selective induction of Mucin 3 by hypoxia in intestinal epithelia. *J. Cell. Biochem.* 99: 1616-1627.

CHROMOSOMAL LOCATION

Genetic locus: Muc3 (mouse) mapping to 5 G2.

PRODUCT

Mucin 3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Mucin 3 shRNA Plasmid (m): sc-149703-SH and Mucin 3 shRNA (m) Lentiviral Particles: sc-149703-V as alternate gene silencing products.

For independent verification of Mucin 3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149703A, sc-149703B and sc-149703C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Mucin 3 siRNA (m) is recommended for the inhibition of Mucin 3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Mucin 3 (M3.1): sc-7315 is recommended as a control antibody for monitoring of Mucin 3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Mucin 3 gene expression knockdown using RT-PCR Primer: Mucin 3 (m)-PR: sc-149703-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.