



NAIF1 siRNA (m): sc-149806

BACKGROUND

NAIF1 (nuclear apoptosis inducing factor 1) is a 327 amino acid protein that localizes to the nucleus and contains two N-terminal nuclear localization signals, one glycine-rich region and one homeodomain-like region. Expressed in a variety of tissues, NAIF1 interacts with C11orf77 and, when overexpressed, functions to halt cell growth and induce apoptosis. NAIF1 exists as multiple alternatively spliced isoforms which are encoded by a gene that maps to human chromosome 9. Chromosome 9 houses over 900 genes and comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and Familial dysautonomia, are both associated with chromosome 9. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

REFERENCES

1. Lv, B., Shi, T., Wang, X., Song, Q., Zhang, Y., Shen, Y., Ma, D. and Lou, Y. 2006. Overexpression of the novel human gene, nuclear apoptosis-inducing factor 1, induces apoptosis. *Int. J. Biochem. Cell Biol.* 38: 671-683.
2. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610673. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Gardiner, J., Barton, D., Marc, J. and Overall, R. 2007. Potential role of tubulin acetylation and microtubule-based protein trafficking in familial dysautonomia. *Traffic* 8: 1145-1149.
4. Temtamy, S.A., Kamel, A.K., Ismail, S., Helmy, N.A., Aglan, M.S., El Gammal, M., El Ruby, M. and Mohamed, A.M. 2007. Phenotypic and cytogenetic spectrum of 9p trisomy. *Genet. Couns.* 18: 29-48.
5. Sinzelle, L., Kapitonov, V.V., Grzela, D.P., Jursch, T., Jurka, J., Izsvák, Z. and Ivics, Z. 2008. Transposition of a reconstructed Harbinger element in human cells and functional homology with two transposon-derived cellular genes. *Proc. Natl. Acad. Sci. USA* 105: 4715-4720.

CHROMOSOMAL LOCATION

Genetic locus: Naif1 (mouse) mapping to 2 B.

PRODUCT

NAIF1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NAIF1 shRNA Plasmid (m): sc-149806-SH and NAIF1 shRNA (m) Lentiviral Particles: sc-149806-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NAIF1 siRNA (m) is recommended for the inhibition of NAIF1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NAIF1 gene expression knockdown using RT-PCR Primer: NAIF1 (m)-PR: sc-149806-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.