

# NARG1L siRNA (m): sc-149833

## BACKGROUND

They cytoplasmic protein NARG1 (NMDA (N-methyl-d-aspartate) receptor-regulated gene 1) interacts with ARD1 or ARD2 to form a complex, which exhibits N-terminal ( $\alpha$ ) acetyltransferase activity. This complex interacts with ribosomal subunits functioning in cotranslational acetylation. During apoptosis, both NARG1 and ARD1 are cleaved by caspases, which results in decreased acetyltransferase activity. Knockdown of NARG1 in HeLa cells leads to apoptosis, indicating that properly functioning NARG1 is essential for cell viability. NARG1 is expressed at high levels in dividing tissues such as bone marrow, testis and embryonal brain and it is overexpressed in papillary thyroid carcinomas. The NARG1-like protein (NARG1L) is an 864 amino acid protein that contains seven TPR repeats and may also be a component of a complex that displays N-terminal acetyltransferase activity.

## REFERENCES

1. Sugiura, N., Patel, R.G. and Corriveau, R.A. 2001. N-methyl-d-aspartate receptors regulate a group of transiently expressed genes in the developing brain. *J. Biol. Chem.* 276: 14257-14263.
2. Fluge, O., Bruland, O., Akslen, L.A., Varhaug, J.E. and Lillehaug, J.R. 2002. NATH, a novel gene overexpressed in papillary thyroid carcinomas. *Oncogene* 21: 5056-5068.
3. Sugiura, N., Adams, S.M. and Corriveau, R.A. 2003. An evolutionarily conserved N-terminal acetyltransferase complex associated with neuronal development. *J. Biol. Chem.* 278: 40113-40120.
4. Asaumi, M., Iijima, K., Sumioka, A., Iijima-Ando, K., Kirino, Y., Nakaya, T. and Suzuki, T. 2005. Interaction of N-terminal acetyltransferase with the cytoplasmic domain of  $\beta$ -amyloid precursor protein and its effect on A $\beta$  secretion. *J. Biochem.* 137: 147-155.
5. Arnesen, T., Anderson, D., Baldersheim, C., Lanotte, M., Varhaug, J.E. and Lillehaug, J.R. 2005. Identification and characterization of the human ARD1-NATH protein acetyltransferase complex. *Biochem. J.* 386: 433-443.
6. Arnesen, T., Gromyko, D., Horvli, O., Fluge, O., Lillehaug, J. and Varhaug, J.E. 2005. Expression of N-acetyl transferase human and human Arrest defective 1 proteins in thyroid neoplasms. *Thyroid* 15: 1131-1136.
7. Arnesen, T., Betts, M.J., Pendino, F., Liberles, D.A., Anderson, D., Caro, J., Kong, X., Varhaug, J.E. and Lillehaug, J.R. 2006. Characterization of hARD2, a processed hARD1 gene duplicate, encoding a human protein N- $\alpha$ -acetyltransferase. *BMC Biochem.* 7: 13-13.
8. Arnesen, T., Gromyko, D., Pendino, F., Rynningen, A., Varhaug, J.E. and Lillehaug, J.R. 2006. Induction of apoptosis in human cells by RNAi-mediated knockdown of hARD1 and NATH, components of the protein N- $\alpha$ -acetyltransferase complex. *Oncogene* 25: 4350-4360.

## CHROMOSOMAL LOCATION

Genetic locus: Naa16 (mouse) mapping to 14 D3.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

NARG1L siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NARG1L shRNA Plasmid (m): sc-149833-SH and NARG1L shRNA (m) Lentiviral Particles: sc-149833-V as alternate gene silencing products.

For independent verification of NARG1L (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149833A, sc-149833B and sc-149833C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

NARG1L siRNA (m) is recommended for the inhibition of NARG1L expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NARG1L gene expression knockdown using RT-PCR Primer: NARG1L (m)-PR: sc-149833-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.