

Nkx-2.4 siRNA (m): sc-150000

BACKGROUND

Members of the NK-2 family of homeodomain proteins, which include Nkx-2.2, Nkx-2.4, Nkx-2.5, Nkx-2.6 and many others, are key regulators of growth and development in several tissues, including brain, heart and pancreas. Nkx-2.2 is responsible for directing ventral neuronal patterning in response to graded Shh signaling. Nkx-2.4 is a 354 amino acid nuclear protein that is detected in the posterior hypothalamus of the embryo and also in adult testis. Nkx-2.5, also designated cardiac specific homeobox protein (Csx), is a homolog of the *Drosophila* tinman protein and is essential for normal cardiovascular development. Nkx-2.6, also a homolog of the *Drosophila* tinman protein, is expressed in the caudal pharyngeal pouches, the caudal heart progenitors, the sinus venosus, the outflow tract of the heart and in a short segment of the gut between stages E8.5 and E10.5 of embryogenesis.

REFERENCES

- Price, M., Lazzaro, D., Pohl, T., Mattei, M.G., Rüther, U., Olivo, J.C., Duboule, D. and Di Lauro, R. 1992. Regional expression of the homeobox gene Nkx-2.2 in the developing mammalian forebrain. *Neuron* 8: 241-255.
- Sussel, L., Kalamaras, J., Hartigan-O'Connor, D.J., Meneses, J.J., Pedersen, R.A., Rubenstein, J.L. and German, M.S. 1998. Mice lacking the homeodomain transcription factor Nkx-2.2 have diabetes due to arrested differentiation of pancreatic β cells. *Development* 125: 2213-2221.
- Pabst, O., Herbrand, H. and Arnold, H.H. 1998. Nkx-2.9 is a novel homeobox transcription factor which demarcates ventral domains in the developing mouse CNS. *Mech. Dev.* 73: 85-93.
- McMahon, A.P. 2000. Neural patterning: the role of Nkx genes in the ventral spinal cord. *Genes Dev.* 14: 2261-2264.
- Wang, C.C., Brodnicki, T., Copeland, N.G., Jenkins, N.A. and Harvey, R.P. 2000. Conserved linkage of NK-2 homeobox gene pairs Nkx2-2/2-4 and Nkx2-1/2-9 in mammals. *Mamm. Genome* 11: 466-468.
- Holland, P.W., Booth, H.A. and Bruford, E.A. 2007. Classification and nomenclature of all human homeobox genes. *BMC Biol.* 5: 47.
- Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 607808. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Nkx2-4 (mouse) mapping to 2 G2.

PRODUCT

Nkx-2.4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Nkx-2.4 shRNA Plasmid (m): sc-150000-SH and Nkx-2.4 shRNA (m) Lentiviral Particles: sc-150000-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Nkx-2.4 siRNA (m) is recommended for the inhibition of Nkx-2.4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Nkx-2.4 gene expression knockdown using RT-PCR Primer: Nkx-2.4 (m)-PR: sc-150000-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.