

NNP-1 siRNA (m): sc-150012

BACKGROUND

Novel nuclear protein 1 (NNP-1), also known as RRP1-like protein or Nucleolar protein Nop52, is a 461 amino acid protein belonging to the RRP1 family. Localized to the nucleolus, NNP-1 has simian virus 40-type and bipartite nuclear localization signals and four coiled-coil domains within its C-terminal region. NNP-1 has been found to play an important role in the generation of 28S rRNA in the late processing steps of ribosome biogenesis. At the end of mitosis, nucleolar proteins assemble in a sequential order during the rebuilding of the nucleolus. NNP-1 assembles after Fibrillarin and C23, and simultaneously with B23 and POP1 in the prenucleolar body pathway.

REFERENCES

1. Jansen, E., Meulemans, S.M., Orlans, I.C. and Van de Ven, W.J. 1997. The NNP-1 gene (D21S2056E), which encodes a novel nuclear protein, maps in close proximity to the cystatin B gene within the EPM1 and APECED critical region on 21q22.3. *Genomics* 42: 336-341.
2. Savino, T.M., Bastos, R., Jansen, E. and Hernandez-Verdun, D. 1999. The nucleolar antigen Nop52, the human homologue of the yeast ribosomal RNA processing RRP1, is recruited at late stages of nucleologenesis. *J. Cell Sci.* 112: 1889-1900.
3. Savino, T.M., Gebrane-Younès, J., De Mey, J., Sibarita, J.B. and Hernandez-Verdun, D. 2001. Nucleolar assembly of the rRNA processing machinery in living cells. *J. Cell Biol.* 153: 1097-1110.
4. Scherl, A., Coute, Y., Deon, C., Calle, A., Kindbeiter, K., Sanchez, J.C., Greco, A., Hochstrasser, D. and Diaz, J.J. 2002. Functional proteomic analysis of human nucleolus. *Mol. Biol. Cell* 13: 4100-4109.
5. Zharskaia, O.O. and Zatsepina, O.V. 2005. Assembly of nucleolus-derived foci in various cultured mammalian cells during mitosis. *Tsitologiya* 47: 780-788.
6. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610653. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Zharskaya, O.O., Barsukova, A.S. and Zatsepina, O.V. 2008. Effect of roscovitine, a selective cyclin B-dependent kinase 1 inhibitor, on assembly of the nucleolus in mitosis. *Biochemistry* 73: 411-419.
8. Muro, E., Hoang, T.Q., Jobart-Malfait, A. and Hernandez-Verdun, D. 2008. In nucleoli, the steady state of nucleolar proteins is leptomycin B-sensitive. *Biol. Cell* 100: 303-313.

CHROMOSOMAL LOCATION

Genetic locus: Rrp1 (mouse) mapping to 10 C1.

PRODUCT

NNP-1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NNP-1 shRNA Plasmid (m): sc-150012-SH and NNP-1 shRNA (m) Lentiviral Particles: sc-150012-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NNP-1 siRNA (m) is recommended for the inhibition of NNP-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

NNP-1 (A-6): sc-398970 is recommended as a control antibody for monitoring of NNP-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NNP-1 gene expression knockdown using RT-PCR Primer: NNP-1 (m)-PR: sc-150012-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.