

ORP-10 siRNA (m): sc-151318

BACKGROUND

Members of the oxysterol-binding protein (OSBP) family function as intracellular lipid receptors. OSBPs are involved in lipid metabolism and signal transduction, as well as vesicle transport, and can translocate to the periphery of Golgi membranes when they are bound to oxysterols. ORPs (OSBP-related proteins) belong to a subfamily of OSBPs and consists of ORP-1 through ORP-11. The ORPs have a highly conserved OSBP-type sterol-binding region and a pleckstrin homology domain. They strongly bind to phosphatidic acid and weakly bind to phosphatidylinositol 3-phosphate. The gene encoding the 764 amino acid ORP-10 (oxysterol-binding protein-related protein 10) protein maps to human chromosome 3p23, which spans 200 million base pairs and encodes between 1,100 and 1,500 genes.

REFERENCES

1. Jaworski, C.J., Moreira, E., Li, A., Lee, R. and Rodriguez, I.R. 2001. A family of 12 human genes containing oxysterol-binding domains. *Genomics* 78: 185-196.
2. Lehto, M., Laitinen, S., Chinetti, G., Johansson, M., Ehnholm, C., Staels, B., Ikonen, E. and Olkkonen, V.M. 2001. The OSBP-related protein family in humans. *J. Lipid Res.* 42: 1203-1213.
3. Annis, A.M., Apostolopoulos, J., Dworkin, S., Purton, L.E. and Sparrow, R.L. 2002. An oxysterol-binding protein family identified in the mouse. *DNA Cell Biol.* 21: 571-580.
4. Lehto, M. and Olkkonen, V.M. 2003. The OSBP-related proteins: a novel protein family involved in vesicle transport, cellular lipid metabolism, and cell signalling. *Biochim. Biophys. Acta* 1631: 1-11.
5. Olkkonen, V.M. and Levine, T.P. 2004. Oxysterol binding proteins: in more than one place at one time? *Biochem. Cell Biol.* 82: 87-98.
6. Skirpan, A.L., Dowd, P.E., Sijacic, P., Jaworski, C.J., Gilroy, S. and Kao, T.H. 2006. Identification and characterization of PiORP1, a *Petunia* oxysterol-binding-protein related protein involved in receptor-kinase mediated signalling in pollen, and analysis of the ORP gene family in *Arabidopsis*. *Plant Mol. Biol.* 61: 553-565.
7. Suchanek, M., Hynynen, R., Wohlfahrt, G., Lehto, M., Johansson, M., Saarinen, H., Radzikowska, A., Thiele, C. and Olkkonen, V.M. 2007. The mammalian oxysterol-binding protein-related proteins (ORPs) bind 25-hydroxycholesterol in an evolutionarily conserved pocket. *Biochem. J.* 405: 473-480.
8. Lessmann, E., Ngo, M., Leitges, M., Minguet, S., Ridgway, N.D. and Huber, M. 2007. Oxysterol-binding protein-related protein (ORP) 9 is a PDK-2 substrate and regulates Akt phosphorylation. *Cell. Signal.* 19: 384-392.
9. Yan, D., Mäyränpää, M.I., Wong, J., Perttilä, J., Lehto, M., Jauhiainen, M., Kovanen, P.T., Ehnholm, C., Brown, A.J. and Olkkonen, V.M. 2008. OSBP-related protein 8 (ORP8) suppresses ABCA1 expression and cholesterol efflux from macrophages. *J. Biol. Chem.* 283: 332-340.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: *Osbpl10* (mouse) mapping to 9 F3.

PRODUCT

ORP-10 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ORP-10 shRNA Plasmid (m): sc-151318-SH and ORP-10 shRNA (m) Lentiviral Particles: sc-151318-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ORP-10 siRNA (m) is recommended for the inhibition of ORP-10 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ORP-10 gene expression knockdown using RT-PCR Primer: ORP-10 (m)-PR: sc-151318-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.