

# Pdcd-4 siRNA (m): sc-152123

## BACKGROUND

The transformation suppressor gene Pdcd-4 (programmed cell death gene 4) inhibits the tumor-promoter-mediated transformation of mouse keratinocytes and is a potential tumor suppressor gene in the development of human lung cancer. Biochemical analysis suggests that the Pdcd-4 protein is involved in protein translation as well as in nuclear events. Pdcd-4 directly interacts with the RNA helicase eIF4A and inhibits protein synthesis by interfering with the assembly of the cap-dependent translation initiation complex. Pdcd-4 also suppresses the transactivation of AP-1-responsive promoters by c-Jun, suggesting that the transformation suppressor activity of Pdcd-4 might be due, at least in part, to the inhibition of c-Jun activity. In addition to affecting c-Jun phosphorylation, Pdcd-4 blocks the recruitment of the co-activator p300 by c-Jun.

## REFERENCES

1. Lankat-Buttgereit, B., et al. 2003. Programmed cell death protein 4 (Pdcd-4): a novel target for antineoplastic therapy? *Biol. Cell* 95: 515-519.
2. Bohm, M., et al. 2003. The transformation suppressor protein Pdcd-4 shuttles between nucleus and cytoplasm and binds RNA. *Oncogene* 22: 4905-4910.
3. Bitomsky N., et al. 2004. Transformation suppressor protein Pdcd-4 interferes with JNK-mediated phosphorylation of c-Jun and recruitment of the coactivator p300 by c-Jun. *Oncogene* 23: 7484-93.
4. Lankat-Buttgereit B., et al. 2004. Pdcd-4 inhibits growth of tumor cells by suppression of carbonic anhydrase type II. *Mol. Cell. Endocrinol.* 214: 149-153.

## CHROMOSOMAL LOCATION

Genetic locus: Pdcd4 (mouse) mapping to 19 D2.

## PRODUCT

Pdcd-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Pdcd-4 shRNA Plasmid (m): sc-152123-SH and Pdcd-4 shRNA (m) Lentiviral Particles: sc-152123-V as alternate gene silencing products.

For independent verification of Pdcd-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152123A, sc-152123B and sc-152123C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Pdcd-4 siRNA (m) is recommended for the inhibition of Pdcd-4 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Pdcd-4 (B-4): sc-376430 is recommended as a control antibody for monitoring of Pdcd-4 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Pdcd-4 gene expression knockdown using RT-PCR Primer: Pdcd-4 (m)-PR: sc-152123-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Su, W., et al. 2017. MicroRNA-21a-5p/PDCD4 axis regulates mesenchymal stem cell-induced neuroprotection in acute glaucoma. *J. Mol. Cell Biol.* 9: 289-301.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.