

PL6 siRNA (m): sc-152286

BACKGROUND

Transmembrane proteins contain transmembrane domains that are usually characterized by α -helical structures. Transmembrane proteins exist as thermodynamically stable hetero- and homodimers that interact with the lipid bilayer and are involved in both material exchange and communication between the cell and the environment. PL6, also referred to as TMEM115 (transmembrane protein 115) or PP6 (placental protein 6), is a 351 amino acid multi-pass membrane protein that is highly expressed in kidney and skeletal muscle, with lower levels of expression detected in liver, placenta, pancreas, lung, heart and brain. PL6 contains one phosphoserine residue and several transmembrane domains, suggesting that it may participate in protein exchange and signaling events between cells.

REFERENCES

1. Popot, J.L. and Engelman, D.M. 1990. Membrane protein folding and oligomerization: the two-stage model. *Biochemistry* 29: 4031-4037.
2. Adamian, L. and Liang, J. 2001. Helix-helix packing and interfacial pairwise interactions of residues in membrane proteins. *J. Mol. Biol.* 311: 891-907.
3. Engelman, D.M., Chen, Y., Chin, C.N., Curran, A.R., Dixon, A.M., Dupuy, A.D., Lee, A.S., Lehnert, U., Matthews, E.E., Reshetnyak, Y.K., Senes, A. and Popot, J.L. 2003. Membrane protein folding: beyond the two stage model. *FEBS Lett.* 555: 122-125.
4. Stevens, T.J., Mizuguchi, K. and Arkin, I.T. 2004. Distinct protein interfaces in transmembrane domains suggest an *in vivo* folding model. *Protein Sci.* 13: 3028-3037.
5. Freeman-Cook, L.L. and Dimaio, D. 2005. Modulation of cell function by small transmembrane proteins modeled on the bovine papillomavirus E5 protein. *Oncogene* 24: 7756-7762.
6. Cao, B., Porollo, A., Adamczak, R., Jarrell, M. and Meller, J. 2006. Enhanced recognition of protein transmembrane domains with prediction-based structural profiles. *Bioinformatics* 22: 303-309.

CHROMOSOMAL LOCATION

Genetic locus: Tmem115 (mouse) mapping to 9 F1.

PRODUCT

PL6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PL6 shRNA Plasmid (m): sc-152286-SH and PL6 shRNA (m) Lentiviral Particles: sc-152286-V as alternate gene silencing products.

For independent verification of PL6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152286A, sc-152286B and sc-152286C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PL6 siRNA (m) is recommended for the inhibition of PL6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PL6 gene expression knockdown using RT-PCR Primer: PL6 (m)-PR: sc-152286-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.