

PLAA siRNA (m): sc-152288

BACKGROUND

WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. PLAA, also known as PLAP (phospholipase A2-activating protein) or DOA1, is a 795 amino acid protein that contains one PFU domain, one PUL domain and seven WD repeats. Via its regulatory domains, PLAA interacts with and activates phospholipase A2 (PLA2), thereby playing an important role in the regulation of inflammatory disease processes.

REFERENCES

- Clark, M.A., et al. 1991. Cloning of a phospholipase A2-activating protein. *Proc. Natl. Acad. Sci. USA* 88: 5418-5422.
- Chopra, A.K., et al. 1999. Molecular characterization of cDNA for phospholipase A2-activating protein. *Biochim. Biophys. Acta* 1444: 125-130.
- Ruiz, A., et al. 1999. Cloning of the human phospholipase A2-activating protein (hPLAP) gene on the chromosome 9p21 melanoma deleted region. *Gene* 239: 155-161.
- Beatty, B.G., et al. 1999. Chromosomal localization of phospholipase A2-activating protein, an Ets2 target gene, to 9p21. *Genomics* 62: 529-532.
- Kozlenkov, A., et al. 2002. Function assignment to conserved residues in mammalian alkaline phosphatases. *J. Biol. Chem.* 277: 22992-22999.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603873. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Koumanov, K., et al. 2003. Bimodal regulatory effect of Melittin and phospholipase A2-activating protein on human type II secretory phospholipase A2. *Cell Biol. Int.* 27: 871-877.
- Schwartz, Z., et al. 2005. Phospholipase A2-activating protein (PLAA) is required for 1 α ,25(OH)₂D₃ signaling in growth plate chondrocytes. *J. Cell. Physiol.* 203: 54-70.

CHROMOSOMAL LOCATION

Genetic locus: Plaa (mouse) mapping to 4 C5.

PRODUCT

PLAA siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PLAA shRNA Plasmid (m): sc-152288-SH and PLAA shRNA (m) Lentiviral Particles: sc-152288-V as alternate gene silencing products.

For independent verification of PLAA (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152288A, sc-152288B and sc-152288C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PLAA siRNA (m) is recommended for the inhibition of PLAA expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PLAA (E-1): sc-390454 is recommended as a control antibody for monitoring of PLAA gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PLAA gene expression knockdown using RT-PCR Primer: PLAA (m)-PR: sc-152288-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.