

PLC η 2 siRNA (m): sc-152297

BACKGROUND

Phosphoinositide-specific phospholipase C (PLC) plays a crucial role in the initiation of receptor mediated signal transduction through the generation of the two second messengers, inositol 1,4,5-triphosphate (IP3) and diacylglycerol (DAG) from phosphatidylinositol 4,5-bisphosphate. There are many mammalian PLC isozymes, including PLC β 1, PLC β 2, PLC β 3, PLC β 4, PLC γ 1, PLC γ 2, PLC δ 1, PLC δ 2, PLC ϵ and PLC η 2. PLC η 2 (phospholipase C, η 2), also known as PLCH2 or PLCL4, is a 1,416 amino acid cell membrane protein that contains one C2 domain, one PH domain, one PI-PLC X-box and Y-box domain and two EF-hand domains. Expressed in kidney and retinal tissue, PLC η 2 uses calcium as a cofactor to produce DAG and IP3, thereby playing an important role in the formation and maintenance of neuronal networks. Multiple isoforms of PLC η 2 exist due to alternative splicing events.

REFERENCES

1. Cefai, D., Debre, P., Kaczorek, M., Idziorek, T., Autran, B. and Bismuth, G. 1990. Human immunodeficiency virus-1 glycoproteins gp120 and gp160 specifically inhibit the CD3/T cell-antigen receptor phosphoinositide transduction pathway. *J. Clin. Invest.* 86: 2117-2124.
2. Zhou, Y., Wing, M.R., Sondek, J. and Harden, T.K. 2005. Molecular cloning and characterization of PLC η 2. *Biochem. J.* 391: 667-676.
3. Dotson, C.D., Roper, S.D. and Spector, A.C. 2005. PLC β 2-independent behavioral avoidance of prototypical bitter-tasting ligands. *Chem. Senses* 30: 593-600.
4. Stewart, A.J., Mukherjee, J., Roberts, S.J., Lester, D. and Farquharson, C. 2005. Identification of a novel class of mammalian phosphoinositide-specific phospholipase C enzymes. *Int. J. Mol. Med.* 15: 117-121.
5. Nakahara, M., Shimozaawa, M., Nakamura, Y., Irino, Y., Morita, M., Kudo, Y. and Fukami, K. 2005. A novel phospholipase C, PLC η 2, is a neuron-specific isozyme. *J. Biol. Chem.* 280: 29128-29134.
6. Zhou, Y., Sondek, J. and Harden, T.K. 2008. Activation of human phospholipase C- η 2 by G β γ . *Biochemistry* 47: 4410-4417.

CHROMOSOMAL LOCATION

Genetic locus: Plch2 (mouse) mapping to 4 E2.

PRODUCT

PLC η 2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PLC η 2 shRNA Plasmid (m): sc-152297-SH and PLC η 2 shRNA (m) Lentiviral Particles: sc-152297-V as alternate gene silencing products.

For independent verification of PLC η 2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152297A, sc-152297B and sc-152297C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PLC η 2 siRNA (m) is recommended for the inhibition of PLC η 2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PLC η 2 gene expression knockdown using RT-PCR Primer: PLC η 2 (m)-PR: sc-152297-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.