

Polr2j siRNA (m): sc-152374

BACKGROUND

RNA polymerase II (Pol II) is a multi-subunit enzyme responsible for the transcription of protein-coding genes. Transcription initiation requires recruitment of the complete transcription machinery to a promoter via solicitation by activators and chromatin remodeling factors. Pol II can coordinate 10 to 14 subunits. This complex interacts with the promoter regions of genes and a variety of elements and transcription factors. POLR2J (DNA-directed RNA polymerase II subunit J) exist as three variants POLR2J1 (RPB11- α), POLR2J2 (RPB11- β 1), and POLR2J3 (RPB11- β 2), of nuclear protein belonging to the eukaryotic RPB11 RNA polymerase subunit family. POLR2J1/2/3 are DNA-dependent RNA polymerases that catalyze the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. They are also part of the core element with the central large cleft.

REFERENCES

1. Bushnell, D.A., et al. 2004. Structural basis of transcription: an RNA polymerase II-TFIIB cocystal at 4.5 Angstroms. *Science* 303: 983-988.
2. Palangat, M., et al. 2004. Downstream DNA selectively affects a paused conformation of human RNA polymerase II. *J. Mol. Biol.* 341: 429-442.
3. Zhong, S., et al. 2004. Epidermal growth factor enhances cellular TATA binding protein levels and induces RNA polymerase I- and III-dependent gene activity. *Mol. Cell. Biol.* 24: 5119-5129.
4. Cabart, P., et al. 2004. BRCA1 cooperates with NUFIP and P-TEF β to activate transcription by RNA polymerase II. *Oncogene* 23: 5316-5329.
5. Shpakovskii, D.G., et al. 2004. New genes on human chromosome 7: bioinformatic analysis of a gene cluster from the POLR2J family. *Bioorg. Khim.* 30: 621-625.
6. Meka, H., et al. 2005. Crystal structure and RNA binding of the Rpb4/Rpb7 subunits of human RNA polymerase II. *Nucleic Acids Res.* 33: 6435-6444.
7. Ujvári, A. and Luse, D.S. 2006. RNA emerging from the active site of RNA polymerase II interacts with the Rpb7 subunit. *Nat. Struct. Mol. Biol.* 13: 49-54.
8. Cocjocar, M., et al. 2007. Genomic location of the human RNA polymerase II general machinery: evidence for a role of TFIIF and Rpb7 at both early and late stages of transcription. *Biochem. J.* 409: 139-147.

CHROMOSOMAL LOCATION

Genetic locus: Polr2j (mouse) mapping to 5 G2.

PRODUCT

Polr2j siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Polr2j shRNA Plasmid (m): sc-152374-SH and Polr2j shRNA (m) Lentiviral Particles: sc-152374-V as alternate gene silencing products.

For independent verification of Polr2j (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152374A and sc-152374B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Polr2j siRNA (m) is recommended for the inhibition of Polr2j expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

POLR2J1/2/3 (G-2): sc-514129 is recommended as a control antibody for monitoring of POLR2J1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Polr2j gene expression knockdown using RT-PCR Primer: Polr2j (m)-PR: sc-152374-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.