

RAPGEFL1 siRNA (m): sc-152704

BACKGROUND

The Ras superfamily of GTPases can be subdivided into the Ras, Rho/Rac, Sar, Rab, Arf, Rap and Ran subfamilies, all of which control multiple aspects of cell function, including cytoskeletal rearrangement, nuclear signaling and cell growth. The Ras superfamily of GTPases function as regulated switches that toggle between a biologically active GTP-bound and an inactive GDP-bound form. This activation is catalyzed by guanine nucleotide exchange factors (GEFs). RAPGEFL1 (Rap guanine nucleotide exchange factor (GEF)-like 1), also known as Link-GEF1, is a 456 amino acid Rap GEF that may function to activate various Rap proteins. Additionally, RAPGEFL1 is thought to play a role in growth-factor signaling cascades and may be involved in neurite formation and growth.

REFERENCES

1. Kawasaki, H., Springett, G.M., Toki, S., Canales, J.J., Harlan, P., Blumenstiel, J.P., Chen, E.J., Bany, I.A., Mochizuki, N., Ashbacher, A., Matsuda, M., Housman, D.E. and Graybiel, A.M. 1998. A Rap guanine nucleotide exchange factor enriched highly in the basal ganglia. *Proc. Natl. Acad. Sci. USA* 95: 13278-13283.
2. Ebinu, J.O., Bottorff, D.A., Chan, E.Y., Stang, S.L., Dunn, R.J. and Stone, J.C. 1998. RasGRP, a Ras guanyl nucleotide-releasing protein with calcium- and diacylglycerol-binding motifs. *Science* 280: 1082-1086.
3. Reid, A.G. and Gribble, S.M. 2005. SmartCapture and the frontiers of FISH technology: report of the digital scientific UK SmartCapture user's meeting, peterhouse college cambridge, UK, 2nd september 2005. *Chromosome Res.* 13: 835-838.
4. Gribble, S.M., Kalaitzopoulos, D., Burford, D.C., Prigmore, E., Selzer, R.R., Ng, B.L., Matthews, N.S., Porter, K.M., Curley, R., Lindsay, S.J., Baptista, J., Richmond, T.A. and Carter, N.P. 2007. Ultra-high resolution array painting facilitates breakpoint sequencing. *J. Med. Genet.* 44: 51-58.
5. Gielen, S.C., Santegoets, L.A., Kühne, L.C., Van Ijcken, W.F., Boers-Sijmons, B., Hanifi-Moghaddam, P., Helmerhorst, T.J., Blok, L.J. and Burger, C.W. 2007. Genomic and nongenomic effects of estrogen signaling in human endometrial cells: involvement of the growth factor receptor signaling downstream AKT pathway. *Reprod. Sci.* 14: 646-654.

CHROMOSOMAL LOCATION

Genetic locus: Rapgef1 (mouse) mapping to 11 D.

PRODUCT

RAPGEFL1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RAPGEFL1 shRNA Plasmid (m): sc-152704-SH and RAPGEFL1 shRNA (m) Lentiviral Particles: sc-152704-V as alternate gene silencing products.

For independent verification of RAPGEFL1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152704A, sc-152704B and sc-152704C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RAPGEFL1 siRNA (m) is recommended for the inhibition of RAPGEFL1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

RAPGEFL1 (R-93): sc-81908 is recommended as a control antibody for monitoring of RAPGEFL1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RAPGEFL1 gene expression knockdown using RT-PCR Primer: RAPGEFL1 (m)-PR: sc-152704-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.