



RASL12 siRNA (m): sc-152714

BACKGROUND

The Ras family of proteins participate in a diverse array of functions, including signal transduction, cytoskeleton dynamics and intracellular trafficking. Ras proteins cycle between a biologically inactive GDP-bound state and an active GTP-bound state. RASL12 (Ras-like protein family member 12), is a 266 amino acid protein that belongs to the small GTPase superfamily and the Ras family. The RASL12 gene is conserved in chimpanzee, canine, bovine, mouse, rat, chicken, zebrafish and *C. elegans*, and maps to human chromosome 15q22.31. Encoding more than 700 genes, chromosome 15 is made up of approximately 106 million base pairs and is about 3% of the human genome. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is located on chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

REFERENCES

1. Bourne, H.R., Sanders, D.A. and McCormick, F. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
2. Boguski, M.S. and McCormick, F. 1993. Proteins regulating Ras and its relatives. *Nature* 366: 643-654.
3. Whitehead, I.P., Khosravi-Far, R., Kirk, H., Trigo-Gonzalez, G., Der, C.J. and Kay, R. 1996. Expression cloning of lsc, a novel oncogene with structural similarities to the Dbl family of guanine nucleotide exchange factors. *J. Biol. Chem.* 271: 18643-18650.
4. Colicelli, J. 2004. Human Ras superfamily proteins and related GTPases. *Sci. STKE* 2004: RE13.
5. Zody, M.C., Garber, M., Sharpe, T., Young, S.K., Rowen, L., O'Neill, K., Whittaker, C.A., Kamal, M., Chang, J.L., Cuomo, C.A., Dewar, K., Fitzgerald, M.G., Kodira, C.D., Madan, A., Qin, S., Yang, X., et al. 2006. Analysis of the DNA sequence and duplication history of human chromosome 15. *Nature* 440: 671-675.
6. Cachón-González, M.B., Wang, S.Z., Lynch, A., Ziegler, R., Cheng, S.H. and Cox, T.M. 2006. Effective gene therapy in an authentic model of Tay-Sachs-related diseases. *Proc. Natl. Acad. Sci. USA* 103: 10373-10378.

CHROMOSOMAL LOCATION

Genetic locus: Rasl12 (mouse) mapping to 9 C.

PRODUCT

RASL12 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RASL12 shRNA Plasmid (m): sc-152714-SH and RASL12 shRNA (m) Lentiviral Particles: sc-152714-V as alternate gene silencing products.

For independent verification of RASL12 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152714A, sc-152714B and sc-152714C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RASL12 siRNA (m) is recommended for the inhibition of RASL12 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RASL12 gene expression knockdown using RT-PCR Primer: RASL12 (m)-PR: sc-152714-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.